

## Exercises Singularity Theory

1. (3 points) Show that  $x^p + y^p$  and  $x^p + y^p + x^2y^2$  ( $p \geq 5$ ) are not right equivalent in  $\mathcal{R}_2$ .
2. (2 points) Let  $f = x^3 + y^3 + z^3 + 3 \cdot c \cdot xyz \in \mathcal{R}_3$ ,  $c \in \mathbb{R}$ ,  $c \neq 0, -1$ .
  - (a) Compute the Milnor number and determinacy of  $f$ .
  - (b) Explain why  $c = -1$  is special.
3. (4 points) Let  $f = x^n$  and  $F(x, t) := f_t(x) := x^n + tx^{n+1}$ .
  - (a) Determine a smooth function  $H(t, x) : I \times (-\epsilon, \epsilon) \rightarrow \mathbb{R}$  (where  $I \subset \mathbb{R}$  is an interval containing 0), such that  $h_t(x) := H(-, t)$  for all  $t \in (-\epsilon, \epsilon)$  is a coordinate change (i.e., a local diffeomorphism at  $0 \in \mathbb{R}$ ), and such that we have  $f \circ h_t = f_t$ .
  - (b) Find a smooth function  $\xi(x, t)$  which satisfies the following equation

$$\xi(x, t) \cdot \partial_x F(X, t) + \partial_t F_t(x, t) = 0$$

- (c) Write down explicitly the following differential equation.

$$\partial_t H(x, t) = \xi(H(X, t), t).$$

- (d) Show that a solution  $H(x, t)$  of the equation  $\partial_t H(x, t) = \xi(H(X, t), t)$  satisfies the identity

$$\frac{dF(H(x, t), t)}{dt} = 0.$$

4. (3 points) Let  $f = x^3 + xy^p \in \mathcal{R}_2$ ,  $p > 2$ . Find the *smallest* integer  $k$  such that  $\mathfrak{m}^k \subset \mathfrak{m}J_f$ .  
(Remark: This calculation is the main step towards establishing the precise value of the determinacy of  $f$ . Namely, one has the following theorem: *If  $f$  is a germ and  $k$  is the smallest integer such that  $\mathfrak{m}^k \subset \mathfrak{m}J_f$  then the determinacy of  $f$  is either  $k$  or  $k - 1$ .*)