

Technische Universität Chemnitz

Sonderforschungsbereich 393

Numerische Simulation auf massiv parallelen Rechnern

Marina Arav Daniel Hershkowitz
Volker Mehrmann Hans Schneider

**The recursive inverse eigenvalue
problem**

Preprint SFB393/99-09

Preprint-Reihe des Chemnitzer SFB 393

SFB393/99-09

March 1999

Abstract

The recursive inverse eigenvalue problem for matrices is studied, where for each leading principle submatrix an eigenvalue and associated left and right eigenvectors are assigned. Existence and uniqueness results as well as explicit formulas are proven, and applications to nonnegative matrices, Z -matrices, M -matrices, symmetric matrices, Stieltjes matrices and inverse M -matrices are considered.

Authors:

Marina Arav

Department of Mathematics,
Technion
Haifa 32000
Israel

Daniel Hershkovitz¹

Department of Mathematics,
Technion
Haifa 32000
Israel

Volker Mehrmann¹

Fakultät für Mathematik,
TU Chemnitz,
D-09107 Chemnitz,
FRG.

Hans Schneider¹

Mathematics Department,
University of Wisconsin,
Madison, WI 53706,
USA.

¹Partial support received through Sonderforschungsbereich 393 *Numerische Simulation auf massiv parallelen Rechnern* at TU Chemnitz.

1 Introduction

Inverse eigenvalue problems are a very important subclass of inverse problems that arise in the context of mathematical modeling and parameter identification. They have been studied extensively in the last 20 years, see e.g. [3, 5, 7, 11, 12, 14] and the references therein. In particular, the inverse eigenvalue problem for nonnegative matrices is still a topic of very active research, since a necessary and sufficient condition for the existence of a nonnegative matrix with a prescribed spectrum is still an open problem, see [4, 11]. In this paper we study inverse eigenvalue problems in a recursive manner, that allows to extend already constructed solutions if further data become available, as is frequently the case in inverse eigenvalue problems, e.g. [3].

We investigate the following *recursive inverse eigenvalue problem of order n* :

Let F be a field, let $s_1, \dots, s_n \in F$ and let

$$l_1 = \begin{bmatrix} l_{1,1} \end{bmatrix}, \quad l_2 = \begin{bmatrix} l_{2,1} \\ l_{2,2} \end{bmatrix}, \quad \dots, \quad l_n = \begin{bmatrix} l_{n,1} \\ \vdots \\ l_{n,n} \end{bmatrix},$$
$$r_1 = \begin{bmatrix} r_{1,1} \end{bmatrix}, \quad r_2 = \begin{bmatrix} r_{1,2} \\ r_{2,2} \end{bmatrix}, \quad \dots, \quad r_n = \begin{bmatrix} r_{1,n} \\ \vdots \\ r_{n,n} \end{bmatrix}$$

be vectors with elements in F . Construct a matrix $A \in F^{n,n}$ such that

$$\begin{cases} l_i^T A \langle i \rangle = s_i l_i^T \\ A \langle i \rangle r_i = s_i r_i \end{cases} \quad i = 1, \dots, n,$$

where $A \langle i \rangle$ denotes the i -th leading principal submatrix of A .

In the sequel we shall use the notation **RIEP**(n) for “the recursive inverse eigenvalue problem of order n ”.

In Section 2 we study the existence and uniqueness of solutions for **RIEP**(n) in the general case. Our main result gives a recursive characterization of the solution for **RIEP**(n). We also obtain a nonrecursive necessary and sufficient condition for unique solvability as well as an explicit formula for the solution in case of uniqueness.

The results of Section 2 are applied in the subsequent sections to special cases. In Section 3 we discuss nonnegative solutions for **RIEP**(n) over the field \mathbb{R} of real numbers. We also introduce a nonrecursive sufficient condition for the existence

of a nonnegative solution for **RIEP**(n). Uniqueness of nonnegative solutions for **RIEP**(n) is discussed in Section 4. In Section 5 we study Z -matrix and M -matrix solutions for **RIEP**(n) over \mathbb{R} . In Section 6 we consider real symmetric solutions for **RIEP**(n) over \mathbb{R} . In Section 7 we consider positive semidefinite real symmetric solutions for **RIEP**(n) over \mathbb{R} . In Section 8 we combine the results of the previous two sections to obtain analogous results for Stieltjes matrices. Finally, in Section 9 we investigate inverse M -matrix solutions for **RIEP**(n).

2 Existence and uniqueness results

In this section we study the existence and uniqueness of solutions for **RIEP**(n) in the general case. For this purpose we introduce some further notation. For the vectors l_i, r_i we set

$$\tilde{l}_i = \begin{bmatrix} l_{i,1} \\ \vdots \\ l_{i,i-1} \end{bmatrix}, \quad \tilde{r}_i = \begin{bmatrix} r_{1,i} \\ \vdots \\ r_{i-1,i} \end{bmatrix}.$$

The case $n = 1$ is easy to verify.

Proposition 1 *If $l_{1,1} = r_{1,1} = 0$ then every 1×1 matrix A solves **RIEP**(1). If either $l_{1,1} \neq 0$ or $r_{1,1} \neq 0$ then $A = [s_1]$ is the unique solution for **RIEP**(1).*

For $n \geq 2$ we have the following recursive characterization of the solution for **RIEP**(n).

Theorem 2 *Let $n \geq 2$. There exists a solution for **RIEP**(n) if and only if there exists a solution B for **RIEP**($n-1$) such that*

$$l_{n,n} = 0 \quad \Longrightarrow \quad \tilde{l}_n^T B = s_n \tilde{l}_n^T, \quad (1)$$

and

$$r_{n,n} = 0 \quad \Longrightarrow \quad B \tilde{r}_n = s_n \tilde{r}_n. \quad (2)$$

*There exists a unique solution for **RIEP**(n) if and only if there exists a unique solution for **RIEP**($n-1$) and $l_{n,n} r_{n,n} \neq 0$.*

Proof. Let A be an $n \times n$ matrix. Partition A as

$$A = \begin{bmatrix} B & y \\ x^T & z \end{bmatrix}, \quad (3)$$

where B is an $(n-1) \times (n-1)$ matrix. Clearly, A solves **RIEP**(n) if and only if B solves **RIEP**($n-1$) and

$$(s_n I_{n-1} - B) \tilde{r}_n = r_{n,n} y, \quad (4)$$

$$\tilde{l}_n^T (s_n I_{n-1} - B) = l_{n,n} x^T, \quad (5)$$

$$x^T \tilde{r}_n + z r_{n,n} = s_n r_{n,n}, \quad (6)$$

$$\tilde{l}_n^T y + z l_{n,n} = s_n l_{n,n}. \quad (7)$$

It thus follows that there exists a solution for **RIEP**(n) if and only if there exists a solution B for **RIEP**($n-1$) such that the equations (4)–(7) (with unknown x , y and z) are solvable. We now show that these equations are solvable if and only if (1) and (2) hold. Distinguish between four cases:

1. $r_{n,n} = 0$, $l_{n,n} \neq 0$. Here (4) is equivalent to (2), (5) is equivalent to

$$x^T = \frac{\tilde{l}_n^T (s_n I_{n-1} - B)}{l_{n,n}}, \quad (8)$$

and (6) then follows from (4). For every $y \in F^{n-1}$ we can find $z \in F$ such that (7) holds.

2. $l_{n,n} = 0$, $r_{n,n} \neq 0$. Here (5) is equivalent to (1), (4) is equivalent to

$$y = \frac{(s_n I_{n-1} - B)\tilde{r}_n}{r_{n,n}}, \quad (9)$$

and (7) then follows from (5). For every $x \in F^{n-1}$ we can find $z \in F$ such that (6) holds.

3. $l_{n,n} = r_{n,n} = 0$. Here (4) is equivalent to (2) and (5) is equivalent to (1). For any $x \in F^{n-1}$ with $x^T \tilde{r}_n = 0$ we have (6), and for any $y \in F^{n-1}$ with $\tilde{l}_n^T y = 0$ we have (7), where z can be chosen arbitrarily.
4. $l_{n,n} \neq 0$, $r_{n,n} \neq 0$. Here (4)–(7) have a unique solution, given by (8), (9) and

$$z = s_n - \frac{\tilde{l}_n^T (s_n I_{n-1} - B)\tilde{r}_n}{l_{n,n} r_{n,n}}. \quad (10)$$

It follows that (4)–(7) are solvable if and only if (1) and (2) hold.

To prove the uniqueness assertion, note that it follows from our proof that if either $l_{n,n} = 0$ or $r_{n,n} = 0$ then a solution is not unique, since at least one of the vectors x , y and z can be chosen arbitrarily. If both $l_{n,n} \neq 0$ and $r_{n,n} \neq 0$ then every solution B for **RIEP**($n-1$) defines a unique solution A for **RIEP**(n). The uniqueness claim follows. \square

This result is recursive and allows to derive a recursive algorithm to compute the solution, but we do not get explicit nonrecursive conditions that characterize the existence of solutions. In order to get a necessary and sufficient condition for unique solvability as well as an explicit formula for the solution in case of uniqueness, we define the $n \times n$ matrix R_n to be the matrix whose columns are r_1, \dots, r_n with zeros appended at the bottom to obtain n -vectors. Similarly, we

define the $n \times n$ matrix L_n to be the matrix whose rows are l_1, \dots, l_n with zeros appended at the right to obtain n -vectors. That is, we have

$$L_n = \begin{bmatrix} l_{1,1} & & & & \\ l_{2,1} & l_{2,2} & & & \\ \vdots & & \ddots & & \\ l_{n,1} & \cdots & l_{n,n-1} & l_{n,n} & \end{bmatrix}, \quad R_n = \begin{bmatrix} r_{1,1} & r_{1,2} & \cdots & r_{1,n} \\ & r_{2,2} & & \vdots \\ & & \ddots & r_{n-1,n} \\ & & & r_{n,n} \end{bmatrix}. \quad (11)$$

We denote

$$S_n = \begin{bmatrix} s_1 & s_2 & s_3 & \cdots & s_n \\ s_2 & s_2 & s_3 & \cdots & s_n \\ s_3 & s_3 & s_3 & \cdots & s_n \\ \vdots & & & & \vdots \\ s_n & s_n & \cdots & \cdots & s_n \end{bmatrix}. \quad (12)$$

Also, we denote by \circ the Hadamard (or elementwise) product of matrices.

Proposition 3 *A solution A for $\mathbf{RIEP}(n)$ satisfies*

$$L_n A R_n = S_n \circ (L_n R_n). \quad (13)$$

Proof. We prove our claim by induction on n . For $n = 1$ the claim follows easily. Assume that the assertion holds for $n < k$ and let $n = k$. Partition A as in (3). We have

$$\begin{aligned} L_n A R_n &= \begin{bmatrix} L_{n-1} & 0 \\ \tilde{l}_n^T & l_{n,n} \end{bmatrix} \begin{bmatrix} B & y \\ x^T & z \end{bmatrix} \begin{bmatrix} R_{n-1} & \tilde{r}_n \\ 0 & r_{n,n} \end{bmatrix} \\ &= \begin{bmatrix} L_{n-1} B R_{n-1} & L_{n-1} (B \tilde{r}_n + r_{n,n} y) \\ (\tilde{l}_n^T B + l_{n,n} x^T) R_{n-1} & (\tilde{l}_n^T B + l_{n,n} x^T) \tilde{r}_n + (\tilde{l}_n^T y + l_{n,n} z) r_{n,n} \end{bmatrix}. \end{aligned}$$

By the inductive assumption we have $L_{n-1} B R_{n-1} = S_{n-1} \circ (L_{n-1} R_{n-1})$. Also, by (4) we have $B \tilde{r}_n + r_{n,n} y = s_n \tilde{r}_n$, by (5) we have $\tilde{l}_n^T B + l_{n,n} x^T = s_n \tilde{l}_n^T$, and by (7) we have $\tilde{l}_n^T y + l_{n,n} z = s_n l_{n,n}$. It thus follows that

$$L_n A R_n = \begin{bmatrix} S_{n-1} \circ (L_{n-1} R_{n-1}) & s_n L_{n-1} \tilde{r}_n \\ s_n \tilde{l}_n^T R_{n-1} & s_n (\tilde{l}_n^T \tilde{r}_n + l_{n,n} r_{n,n}) \end{bmatrix} = S_n \circ (L_n R_n). \quad \square$$

In general, the converse of Proposition 3 does not hold, that is, a matrix A satisfying (13) does not necessarily form a solution for $\mathbf{RIEP}(n)$, as is demonstrated by Example 5 below.

Theorem 4 *There is a unique solution for $\mathbf{RIEP}(n)$ if and only if*

$$l_{1,1} \neq 0 \quad \text{or} \quad r_{1,1} \neq 0$$

and

$$l_{i,i}r_{i,i} \neq 0, \quad i = 1, \dots, n.$$

Furthermore, the unique solution is given by

$$A = L_n^{-1}[S_n \circ (L_n R_n)]R_n^{-1}. \quad (14)$$

Proof. The uniqueness claim follows from Proposition 1 and Theorem 2. The fact that the unique solution for **RIEP**(n) is given by (14) follows immediately from Proposition 3. \square

In the case that the solution is not unique, that is, whenever $l_{1,1} = r_{1,1} = 0$ or whenever $l_{i,i}$ or $r_{i,i}$ vanish for some $i > 1$, the matrices L_n and R_n defined in (11) are not invertible. Therefore, in this case (14) is invalid. We conclude this section by an example showing that, in general, a revised form of (14), with inverses replaced by generalized inverses, does not provide a solution for **RIEP**(n).

Example 5 Let

$$s_1 = 1, \quad s_2 = 2, \quad s_3 = 3,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}.$$

We have

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}, \quad R = L^T, \quad S = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 2 & 3 \\ 3 & 3 & 3 \end{bmatrix}.$$

Let L^+ and R^+ be the Moore-Penrose inverses of L and R respectively, see [1]. We have

$$A = L^+[S \circ (LR)]R^+ = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1.5 & 1.5 \\ 0 & 1.5 & 1.5 \end{bmatrix}.$$

Since $A \langle 2 \rangle$ does not have an eigenvalue 2, A is not a solution for **RIEP**(3). Note that we still have $L_n A R_n = S_n \circ (L_n R_n)$.

In this section we have characterized solvability of **RIEP**(n) over a general field F in terms of recursive conditions. We have also given a necessary and sufficient condition for unique solvability and an explicit formula for the unique solution. In the following sections we shall discuss the special cases of nonnegative matrices, Z -matrices, M -matrices, real symmetric matrices, positive semidefinite matrices, Stieltjes matrices and inverse M -matrices.

3 Existence of nonnegative solutions

In this section we apply the results of Section 2 to nonnegative solutions for **RIEP**(n) over the field \mathbb{R} of real numbers. A matrix $A \in \mathbb{R}^{n,n}$ is said to be *nonnegative* [*positive*] if all elements of A are nonnegative [*positive*]. In this case we write $A \geq 0$ [$A > 0$]. For matrices $A, B \in \mathbb{R}^{n,n}$ we write $A \geq B$ if $A - B \geq 0$ and $A > B$ if $A - B > 0$.

In order to state our results we define a vector over \mathbb{R} to be *unisign* if its nonzero components have the same sign.

Theorem 6 *Let $n \geq 2$. There exists a nonnegative solution for **RIEP**(n) if and only if we have*

$$l_i \text{ or } r_i \text{ is a unisign nonzero vector} \implies s_i \geq 0, \quad i = 1, \dots, n, \quad (15)$$

and there exists a nonnegative solution B for **RIEP**($n-1$) satisfying

$$\begin{cases} \frac{s_n \tilde{r}_n}{r_{n,n}} \geq \frac{B \tilde{r}_n}{r_{n,n}}, & r_{n,n} \neq 0 \\ s_n \tilde{r}_n = B \tilde{r}_n, & r_{n,n} = 0 \end{cases}, \quad (16)$$

$$\begin{cases} \frac{\tilde{l}_n^T s_n}{l_{n,n}} \geq \frac{\tilde{l}_n^T B}{l_{n,n}}, & l_{n,n} \neq 0 \\ \tilde{l}_n^T s_n = \tilde{l}_n^T B, & l_{n,n} = 0 \end{cases}, \quad (17)$$

and

$$l_{n,n} r_{n,n} \neq 0 \implies s_n \left(\frac{\tilde{l}_n^T \tilde{r}_n}{l_{n,n} r_{n,n}} - 1 \right) \leq \frac{\tilde{l}_n^T B \tilde{r}_n}{l_{n,n} r_{n,n}}. \quad (18)$$

*There exists a positive solution for **RIEP**(n) if and only if there exists a positive solution B for **RIEP**($n-1$) such that (15)–(18) hold with strict inequalities and every nonzero unisign vector l_i or r_i has no zero components.*

Proof. Let $A \in \mathbb{R}^{n,n}$. As in the proof of Theorem 2, partition A as in (3), and so A solves **RIEP**(n) if and only if B solves **RIEP**($n-1$) and (4)–(7) hold. Therefore, if A is a nonnegative solution for **RIEP**(n) then we have (16)–(18). Also, it follows from the nonnegativity of A that (15) holds. Conversely, assume that (15) holds and that B forms a nonnegative solution for **RIEP**($n-1$) satisfying (16)–(18). We show that in this case we can find nonnegative solutions x , y and z for (4)–(7). Distinguish between four cases:

1. $r_{n,n} = 0$, $l_{n,n} \neq 0$. Here x is given by (8), y can be chosen arbitrarily, and z should be chosen such that (7) holds. It follows from (17) that x is nonnegative. If $s_n \geq 0$ then we choose $y = 0$, we have $z = s_n$, and so we have a nonnegative solution for (4)–(7). If $s_n < 0$ then, by (15), l_n is not

unisign and hence $\frac{\tilde{l}_n^T}{l_{n,n}}$ has at least one negative component. It follows that we can find a positive vector y such that $\frac{\tilde{l}_n^T y}{l_{n,n}} < s_n$. Since by (7) we have $z = s_n - \frac{\tilde{l}_n^T y}{l_{n,n}}$, it follows that $z > 0$, and so again we have a nonnegative solution for (4)–(7).

2. $l_{n,n} = 0, r_{n,n} \neq 0$. Here y is given by (9), x can be chosen arbitrarily, and z should be chosen such that (6) holds. The proof follows as in the previous case.
3. $l_{n,n} = r_{n,n} = 0$. Here x and y should be chosen such that $x^T \tilde{r}_n = \tilde{l}_n^T y = 0$ and z can be chosen arbitrarily. In order to obtain a nonnegative solution we can choose x, y and z to be zero.
4. $l_{n,n} \neq 0, r_{n,n} \neq 0$. Here x is given by (8), y is given by (9), and z is given by (10). It follows from (17), (16) and (18) that x, y and z are nonnegative.

Assume now that A is a *positive* solution for **RIEP**(n). It is easy to verify that in this case (15)–(18) should hold with strict inequalities. Also, for every nonzero unisign vector l_i [r_i], the vector $l_i^T A \langle i \rangle$ [$A \langle i \rangle r_i$] has no zero components, implying that $l_i, [r_i]$ has no zero components. Conversely, assume that (15) holds with a strict inequality, that every nonzero unisign vector l_i or r_i has no zero components, and that B forms a positive solution for **RIEP**($n-1$) satisfying (16)–(18) with strict inequalities. We show that in this case we can find positive solutions x, y and z for (4)–(7). Note that in Case 1 above, the vector x now becomes positive. Also, since the inequality in (15) is now strict, we have either $s_n > 0$, in which case we can choose *positive* y sufficiently small such that z is positive, or $s_n \leq 0$, in which case y can be chosen positive as before and the resulting z is positive. The same arguments hold for Case 2. In Case 4, it follows from the strict inequalities (17)–(18) that x, y and z are positive. Finally, in Case 3, since l_n and r_n both have at least one zero component, it follows that both vectors are not unisign. Hence, we can find positive x and y such that $x^T \tilde{r}_n = \tilde{l}_n^T y = 0$. We assign any positive number to z to find a positive solution A for **RIEP**(n). \square

By the Perron-Frobenius theory, see e.g. [8, 2], the largest absolute value $\rho(A)$ of an eigenvalue of a nonnegative $n \times n$ matrix A is itself an eigenvalue of A , the so called *Perron root* of A , and it has an associated nonnegative eigenvector. Furthermore, if A is *irreducible*, that is, if either $n = 1$ or $n \geq 2$ and there exists no permutation matrix P such that $P^T A P = \begin{bmatrix} B & C \\ 0 & D \end{bmatrix}$ where B and D are square, then $\rho(A)$ is a simple eigenvalue of A with an associated positive eigenvector. If A is not necessarily irreducible then we have the following, see e.g. [2].

Theorem 7 *If B is a principal submatrix of a nonnegative square matrix A then $\rho(B) \leq \rho(A)$. Furthermore, $\rho(A)$ is an eigenvalue of some proper principal submatrix of A if and only if A is reducible.*

Note that if we require that the s_i are the Perron roots of the principal submatrices $A\langle i \rangle$, $i = 1, \dots, n$, then, by Theorem 7, we have

$$0 \leq s_1 \leq s_2 \leq \dots \leq s_n. \quad (19)$$

If, furthermore, all the leading principal submatrices of A are required to be irreducible, then

$$0 \leq s_1 < s_2 < \dots < s_n. \quad (20)$$

Condition (19) is not sufficient to guarantee that a nonnegative solution A for **RIEP**(n) necessarily has s_1, \dots, s_n as Perron roots of $A\langle i \rangle$, $i = 1, \dots, n$, as is demonstrated by the following example.

Example 8 Let

$$s_1 = s_2 = 1, \quad s_3 = 2,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}.$$

The nonnegative matrix

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

solves **RIEP**(3). Note that $\rho(A) = 3 > s_3$.

In order to see cases in which s_1, \dots, s_n are the Perron roots of $A\langle i \rangle$, $i = 1, \dots, n$, respectively, we prove

Proposition 9 *If the vector l_n or r_n is positive then for a nonnegative solution A for **RIEP**(n) we have $\rho(A) = s_n$.*

Proof. The claim follows immediately from the known fact that a positive eigenvector of a nonnegative matrix corresponds to the spectral radius, see e.g. Theorem 2.1.11 in [2, p. 128]. \square

Corollary 10 *If for every $i \in \{1, \dots, n\}$ we have either $l_i > 0$ or $r_i > 0$ then for every nonnegative solution A for **RIEP**(n) we have $\rho(A\langle i \rangle) = s_i$, $i = 1, \dots, n$.*

Lemma 11 *Assume that there exists a nonnegative solution A for **RIEP**(n) such that $\rho(A\langle n-1 \rangle) < s_n$. If $r_n \neq 0$ or $l_n \neq 0$ then $\rho(A) = s_n$.*

Proof. Since $r_n \neq 0$ or $l_n \neq 0$ it follows that s_n is an eigenvalue of A . Assume that $s_n \neq \rho(A)$. It follows that the nonnegative matrix A has at least two eigenvalues larger than or equal to s_n . By [6, p. 473], see also [10, Corollary 1], it follows that $\rho(A\langle n-1 \rangle) \geq s_n$, which is a contradiction. Therefore, we have $s_n = \rho(A)$. \square

Corollary 12 *If for every $i \in \{1, \dots, n\}$ we have either $r_i \neq 0$ or $l_i \neq 0$, and if (20) holds then for every nonnegative solution A for **RIEP**(n) we have $\rho(A\langle i \rangle) = s_i$, $i = 1, \dots, n$.*

Proof. Note that $A\langle 1 \rangle = [s_1]$ and so $\rho(A\langle 1 \rangle) = s_1$. Our result follows using Lemma 11 repeatedly. \square

Lemma 13 *Assume that $r_n \geq 0$ and $r_{n,n} \neq 0$ or that $l_n \geq 0$ and $l_{n,n} \neq 0$. Then for every nonnegative solution A for **RIEP**(n) we have $\rho(A) = \max\{\rho(A\langle n-1 \rangle), s_n\}$.*

Proof. Without loss of generality, we consider the case where $r_n \geq 0$ and $r_{n,n} \neq 0$. If r_n is positive then, by Proposition 9, we have $\rho(A) = s_n$ and, since by the Perron-Frobenius theory we have $\rho(A\langle n-1 \rangle) \leq \rho(A)$, the result follows. Otherwise, r_n has some zero components. Let α be the set of indices i such that $r_{i,n} > 0$ and let α^c be the complement of α in $\{1, \dots, n\}$. Note that since r_n is a nonnegative eigenvector of the nonnegative matrix A it follows that the submatrix $A[\alpha^c|\alpha]$ of A , with rows indexed by α^c and columns indexed by α , is a zero matrix. It follows that A is a reducible matrix and $\rho(A) = \max\{\rho(A[\alpha^c|\alpha^c]), \rho(A[\alpha|\alpha])\}$. Note that the subvector $r_n[\alpha]$ of r_n indexed by α is a positive eigenvector of $A[\alpha|\alpha]$ associated with the eigenvalue s_n . It thus follows that $\rho(A[\alpha|\alpha]) = s_n$. Since $n \in \alpha$ it follows that $A[\alpha^c|\alpha^c]$ is a submatrix of $A\langle n-1 \rangle$. Thus, by the Perron-Frobenius theory we have $\rho(A[\alpha^c|\alpha^c]) \leq \rho(A\langle n-1 \rangle) \leq \rho(A)$. Hence, it follows that $\rho(A) = \max\{s_{n-1}, s_n\}$. \square

Corollary 14 *Assume that for every $i \in \{1, \dots, n\}$ we have either $r_i \geq 0$ and $r_{i,i} \neq 0$ or $l_i \geq 0$ and $l_{i,i} \neq 0$. Then for every nonnegative solution A for **RIEP**(n) we have $\rho(A\langle i \rangle) = \max_{j=1, \dots, i}\{s_j\}$.*

Proof. Note that $A\langle 1 \rangle = [s_1]$ and so $\rho(A\langle 1 \rangle) = s_1$. Our result follows using Lemma 13 repeatedly. \square

Corollary 15 *Assume that for every $i \in \{1, \dots, n\}$, we have either $r_i \geq 0$ and $r_{i,i} \neq 0$ or $l_i \geq 0$ and $l_{i,i} \neq 0$. If (19) holds then for every nonnegative solution A we have $\rho(A\langle i \rangle) = s_i$, $i = 1, \dots, n$.*

Another interesting consequence of Theorem 4 is the following relationship between the matrix elements and the eigenvectors associated with the Perron roots of the leading principal submatrices of a nonnegative matrix.

Corollary 16 Let $n \geq 2$. Let $A \in \mathbf{R}^{n,n}$ be a nonnegative matrix, let s_i, l_i and r_i be the Perron roots and associated left and right eigenvectors of $A\langle i \rangle$, $i = 1, \dots, n$, respectively, and assume that (20) holds. Let S_n, L_n, R_n be defined as in (11) and (12). Then

$$A\langle i \rangle = L_i^{-1}[S_i \circ (L_i R_i)]R_i^{-1}, \quad i = 1, \dots, n. \quad (21)$$

Proof. Since (20) holds, it follows that s_i is not an eigenvalue of $A\langle i-1 \rangle$, $i = 2, \dots, n$. Therefore, it follows from (1) and (2) that $l_{i,i}r_{i,i} \neq 0$. Also, since l_1 and r_1 are eigenvectors of $A\langle 1 \rangle$, we have $l_{1,1}r_{1,1} \neq 0$. It now follows from Theorem 4 that $A\langle i \rangle$ is the unique solution for **RIEP**(i), and is given by (21). \square

While Theorem 6 provides a recursive characterization for nonnegative solvability of **RIEP**(n), in general nonrecursive necessary and sufficient conditions for the existence of nonnegative solution are not known. We now present a nonrecursive sufficient condition.

Corollary 17 Assume that the vectors l_i, r_i , $i = 1, \dots, n$ are all positive and that the numbers s_1, \dots, s_n are all positive. Let

$$M_i^r = \max_{j=1, \dots, i-1} \frac{r_{j,i}}{r_{j,i-1}}, \quad m_i^r = \min_{j=1, \dots, i-1} \frac{r_{j,i}}{r_{j,i-1}},$$

$$M_i^l = \max_{j=1, \dots, i-1} \frac{l_{i,j}}{l_{i-1,j}}, \quad m_i^l = \min_{j=1, \dots, i-1} \frac{l_{i,j}}{l_{i-1,j}}.$$

If we have

$$s_i m_i^r \geq s_{i-1} M_i^r, \quad i = 2, \dots, n, \quad (22)$$

$$s_i m_i^l \geq s_{i-1} M_i^l, \quad i = 2, \dots, n, \quad (23)$$

and

$$s_i (\tilde{l}_i^T \tilde{r}_i - l_{i,i} r_{i,i}) \leq s_{i-1} \max \{ m_i^r \tilde{l}_i^T r_{i-1}, m_i^l l_{i-1}^T \tilde{r}_i \}, \quad i = 2, \dots, n, \quad (24)$$

then there exists a (unique) nonnegative solution A for **RIEP**(n).

Furthermore, if all the inequalities (22)-(24) hold with strict inequality then there exists a (unique) positive solution A for **RIEP**(n).

Proof. We prove our assertion by induction on n . The case $n = 1$ is trivial. By the inductive assumption we can find a nonnegative solution B for **RIEP**($n-1$). Note that

$$M_n^r r_{n-1} \geq \tilde{r}_n \geq m_n^r r_{n-1}. \quad (25)$$

Therefore, it follows from (22) that

$$s_n \tilde{r}_n \geq s_n m_n^r r_{n-1} \geq s_{n-1} M_n^r r_{n-1} = M_n^r B r_{n-1} \geq B \tilde{r}_n,$$

and so (16) holds. Similarly we prove that (17) holds. To prove that (18) holds note that by (25) we have $B \tilde{r}_n \geq B m_n^r r_{n-1} = s_{n-1} m_n^r r_{n-1}$.

Similarly, we have $\tilde{l}_n^T B \geq s_{n-1} m_n^l l_{n-1}^T$. Hence, it follows that $\tilde{l}_n^T B \tilde{r}_n \geq s_{n-1} \max \{m_n^r \tilde{l}_n^T r_{n-1}, m_n^l l_{n-1}^T \tilde{r}_n\}$. By (24) applied to $i = n$ we obtain (18). By Theorem 6, there exists a nonnegative solution for **RIEP**(n). The proof of the positive case is similar. \square

The conditions in Corollary 17 are not necessary as is demonstrated by the following example.

Example 18 Let $s_1 = 1$, $s_2 = 2$, $s_3 = 3$ and let

$$r_1 = l_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad r_2 = l_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad r_3 = \begin{bmatrix} 3 \\ 5 \\ 1 \end{bmatrix}, \quad l_3 = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}.$$

We have $m_3^r = 3$, $M_3^r = 5$, $m_3^l = 1$ and $M_3^l = 2$. Note that both (22) and (23) do not hold for $i = 3$. Nevertheless, the unique solution for **RIEP**(3) is the nonnegative matrix

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 7 \\ 1 & 0 & 0 \end{bmatrix}.$$

4 Uniqueness of nonnegative solutions

When considering uniqueness of nonnegative solutions for **RIEP**(n), observe that it is possible that **RIEP**(n) does not have a unique solution but does have a unique nonnegative solution, as is demonstrated by the following example.

Example 19 Let

$$s_1 = s_2 = 0,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad r_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

By Theorem 2, there is no unique solution for **RIEP**(2). Indeed, the solutions for **RIEP**(2) are all matrices of the form

$$\begin{bmatrix} 0 & 0 \\ a & -a \end{bmatrix}.$$

Clearly, the zero matrix is the only nonnegative solution for **RIEP**(2).

Observe that, unlike in Theorem 2, the existence of a unique nonnegative solution for **RIEP**(n) does not necessarily imply the existence of a unique nonnegative solution for **RIEP**($n-1$), as is demonstrated by the following example.

Example 20 Let

$$s_1 = s_2 = 0, \quad s_3 = 2,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad r_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Observe that all matrices of the form

$$\begin{bmatrix} 0 & 0 \\ a & a \end{bmatrix}$$

solve **RIEP**(2), and hence there is no unique nonnegative solution for **RIEP**(2). However, the only nonnegative solution for **RIEP**(3) is the matrix

$$\begin{bmatrix} 0 & 0 & 2 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}.$$

We remark that one can easily produce a similar example with nonnegative vectors r_i and l_i , $i = 1, \dots, n$.

In order to introduce necessary conditions and sufficient conditions for uniqueness of nonnegative solutions for **RIEP**(n) we prove

Lemma 21 *Let $n \geq 2$, and assume that B forms a nonnegative solution for **RIEP**($n-1$) satisfying (15)–(18). Then there exist unique nonnegative vectors x , y and z such that the matrix $\begin{bmatrix} B & y \\ x^T & z \end{bmatrix}$ solves **RIEP**(n) if and only if either $l_{n,n}r_{n,n} \neq 0$, or $s_n = 0$ and l_n is a unisign vector with no zero components, or $s_n = 0$ and r_n is a unisign vector with no zero components.*

Proof. We follow the proof of Theorem 6. Consider the four cases in that proof. In Case 1, the vector x is uniquely determined and any nonnegative assignment for y is valid as long as $z = s_n - \frac{\tilde{l}_n^T y}{l_{n,n}} \geq 0$. If $s_n > 0$ then every nonnegative vector y sufficiently small will do. If $s_n < 0$ then, as is shown in the proof of Theorem 6, we can find a positive y such that $z > 0$, and by continuity arguments there exist infinitely many such vectors y . If $s_n = 0$ then a unique such y exists if and only if there exists a unique nonnegative vector y such that $\frac{\tilde{l}_n^T y}{l_{n,n}} \leq 0$. Clearly, if \tilde{l}_n has a nonpositive component then every vector y whose corresponding component is positive and all other components are zero solves the problem. On the other hand, if $\tilde{l}_n > 0$, which is equivalent to saying that l_n is a unisign vector with no zero components, then the only nonnegative vector y that solves the problem is $y = 0$. Similarly, we prove that, in case 2, a unique nonnegative solution exists

if and only if $s_n = 0$ and r_n is a unisign vector with no zero components. We do not have uniqueness in Case 3 since then z can be chosen arbitrarily. Finally, there is always uniqueness in Case 4. \square

Lemma 21 yields sufficient conditions and necessary conditions for uniqueness of nonnegative solutions for **RIEP**(n). First, observe that if $s_n = 0$ and l_n is a unisign vector with no zero components, or if $s_n = 0$ and r_n is a unisign vector with no zero components, then the zero matrix is the only nonnegative solution of the problem. A less trivial sufficient condition is the following.

Corollary 22 *Let $n \geq 2$, and let A be a nonnegative solution for **RIEP**(n). If $A\langle n-1 \rangle$ forms a unique nonnegative solution for **RIEP**($n-1$) and if $l_{n,n}r_{n,n} \neq 0$, then A is the unique nonnegative solution for **RIEP**(n).*

Necessary conditions are given by the following

Corollary 23 *Let $n \geq 2$. If there exists a unique nonnegative solution for **RIEP**(n) then either $l_{n,n}r_{n,n} \neq 0$, or $s_n = 0$ and l_n is a unisign vector with no zero components, or $s_n = 0$ and r_n is a unisign vector with no zero components.*

The condition $l_{n,n}r_{n,n} \neq 0$ is not sufficient for the uniqueness of a nonnegative solution for **RIEP**(n), as is shown in the following example.

Example 24 Let

$$s_1 = s_2 = s_3 = 0,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad r_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}.$$

Although we have $l_{n,n}r_{n,n} \neq 0$, all matrices of the form

$$\begin{bmatrix} 0 & 0 & 0 \\ a & a & 0 \\ a & a & 0 \end{bmatrix}$$

solve **RIEP**(3), and hence there is no unique nonnegative solution for **RIEP**(3).

5 The Z -matrix and M -matrix case

A real square matrix A is said to be a Z -matrix if it has nonpositive off-diagonal elements. Note that A can be written as $A = \alpha I - B$ where α is a real number and B is a nonnegative matrix. If we further have that $\alpha \geq \rho(B)$ then we say that A is an M -matrix.

In this section we discuss Z -matrix and M -matrix solutions for $\mathbf{RIEP}(n)$ over the field \mathbf{R} of real numbers. The proofs of the results are very similar to the proofs of the corresponding results in Sections 3 and 4 and, thus, are omitted in most cases.

Theorem 25 *Let $n \geq 2$. There exists a Z -matrix solution for $\mathbf{RIEP}(n)$ if and only if there exists a Z -matrix solution B for $\mathbf{RIEP}(n-1)$ satisfying*

$$\begin{cases} \frac{s_n \tilde{r}_n}{r_{n,n}} \leq \frac{B \tilde{r}_n}{r_{n,n}}, & r_{n,n} \neq 0 \\ s_n \tilde{r}_n = B \tilde{r}_n, & r_{n,n} = 0 \end{cases},$$

and

$$\begin{cases} \frac{\tilde{l}_n^T s_n}{l_{n,n}} \leq \frac{\tilde{l}_n^T B}{l_{n,n}}, & l_{n,n} \neq 0 \\ \tilde{l}_n^T s_n = \tilde{l}_n^T B, & l_{n,n} = 0 \end{cases}.$$

Furthermore, if l_n or r_n is positive then a Z -matrix solution for $\mathbf{RIEP}(n)$ is an M -matrix if and only if $s_n \geq 0$.

Proof. The proof of the first part of the theorem is similar to the proof of Theorem 6, observing that here the vectors x and y are required to be nonnegative and that the sign of z is immaterial. The proof of the second part of the Theorem follows, similarly to Proposition 9, from the known fact that a positive eigenvector of a Z -matrix corresponds to the least real eigenvalue. \square

Theorem 26 *Let $n \geq 2$. Let $A \in \mathbf{R}^{n,n}$ be a Z -matrix, let s_i, l_i and r_i be the least real eigenvalues and the corresponding left and right eigenvectors of $A\langle i \rangle$, $i = 1, \dots, n$, respectively, and assume that*

$$s_1 > s_2 > \dots > s_n.$$

Let S_n, L_n, R_n be defined as in (11) and (12). Then

$$A\langle i \rangle = L_i^{-1}[S_i \circ (L_i R_i)]R_i^{-1}, \quad i = 1, \dots, n.$$

For the numbers M_i^r, m_i^r, M_i^l and m_i^l , defined in Corollary 17, we have

Theorem 27 *Assume that the vectors l_i, r_i , $i = 1, \dots, n$ are all positive and that the numbers s_1, \dots, s_n are all positive. If we have*

$$s_i M_i^r \leq s_{i-1} m_i^r, \quad i = 2, \dots, n$$

and

$$s_i M_i^l \leq s_{i-1} m_i^l, \quad i = 2, \dots, n,$$

then there exists a (unique) M -matrix solution A for $\mathbf{RIEP}(n)$.

Theorem 28 *Let $n \geq 2$, let A be a Z -matrix solution for $\mathbf{RIEP}(n)$ and assume that $A\langle n-1 \rangle$ forms a unique Z -matrix solution for $\mathbf{RIEP}(n-1)$. Then A is the unique Z -matrix solution for $\mathbf{RIEP}(n)$ if and only if $l_{n,n}r_{n,n} \neq 0$.*

Here too, unlike in Theorem 2, the existence of a unique Z -matrix solution for $\mathbf{RIEP}(n)$ does not necessarily imply the existence of a unique Z -matrix solution for $\mathbf{RIEP}(n-1)$, as is demonstrated by the following example.

Example 29 Let $s_1 = s_2 = s_3 = 0$, and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad r_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Observe that all matrices of the form

$$\begin{bmatrix} 0 & 0 \\ a & -a \end{bmatrix}$$

solve $\mathbf{RIEP}(2)$, and hence there is no unique Z -matrix solution for $\mathbf{RIEP}(2)$. However, it is easy to verify that the zero matrix is the only Z -matrix solution for $\mathbf{RIEP}(3)$.

6 The real symmetric case

The inverse eigenvalue problem for real symmetric matrices is well studied, see e.g. [3]. In this section we consider symmetric solutions for $\mathbf{RIEP}(n)$ over the field \mathbb{R} of real numbers. We obtain the following consequence of Theorem 2, characterizing the real symmetric case.

Theorem 30 *Let $n \geq 2$. There exists a symmetric solution for $\mathbf{RIEP}(n)$ if and only if there exists a symmetric solution B for $\mathbf{RIEP}(n-1)$ such that the implications (1) and (2) hold, and*

$$l_{n,n}r_{n,n} \neq 0 \quad \implies \quad (s_n I_{n-1} - B) \left(\frac{\tilde{l}_n}{l_{n,n}} - \frac{\tilde{r}_n}{r_{n,n}} \right) = 0. \quad (26)$$

Furthermore, if there exists a unique symmetric solution for $\mathbf{RIEP}(n)$ then $l_{n,n} \neq 0$ or $r_{n,n} \neq 0$.

Proof. Let $A \in \mathbb{R}^{n,n}$. Partition A as in (3), and so A solves $\mathbf{RIEP}(n)$ if and only if B solves $\mathbf{RIEP}(n-1)$ and (4)–(7) hold. It was shown in the proof of Theorem 2 that (4)–(7) are solvable if and only if (1) and (2) hold. Therefore, all we have to show that if B is symmetric then we can find solutions x , y and z for (4)–(7) such that $y = x$ if and only if (26) holds. We go along the four cases discussed in

Theorem 2. In Case 1, the vector x is uniquely determined and the vector y can be chosen arbitrarily. Therefore, in this case we set $y = x$, and z is then uniquely determined. In Case 2, the vector y is uniquely determined and the vector x can be chosen arbitrarily. Thus, in this case we set $x = y$, and z is then uniquely determined. In Case 3, we can choose any x and y as long as $x^T \tilde{r}_n = 0$ and $\tilde{l}_n^T y = 0$. In particular, we can choose $x = y = 0$. Furthermore, z can be chosen arbitrarily. Finally, in Case 4, we have $x = y$ if and only if (26) holds. Note that this is the only case in which, under the requirement that $y = x$, the vectors x , y and z are uniquely determined. \square

We remark that, unlike in Theorem 2, the existence of a unique symmetric solution for **RIEP**(n) does not necessarily imply the existence of a unique symmetric solution for **RIEP**($n-1$), as is demonstrated by the following example.

Example 31 Let

$$s_1 = 1, \quad s_2 = 2, \quad s_3 = 0,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix},$$

$$l_4 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \quad r_4 = \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}.$$

It is easy to verify that all symmetric matrices of the form

$$\begin{bmatrix} 1 & 1 & a \\ 1 & 1 & a \\ a & a & b \end{bmatrix}, \quad a, b \in \mathbb{R}$$

solve **RIEP**(3), while the unique solution for **RIEP**(4) is

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

This example also shows that there may exist a unique solution for **RIEP**(n) even if $l_{i,i} = r_{i,i} = 0$ for some $i \in 1, \dots, n$.

Naturally, although not necessarily, one may expect in the symmetric case to have the condition

$$r_i = l_i, \quad i = 1, \dots, n. \quad (27)$$

Indeed, in this case we have the following corollary of Theorems 2 and 30.

Corollary 32 *Let $n \geq 2$ and assume that (27) holds. The following are equivalent:*

- (i) *There exists a symmetric solution for $\mathbf{RIEP}(n)$.*
- (ii) *There exists a solution for $\mathbf{RIEP}(n)$.*
- (iii) *There exists a symmetric solution B for $\mathbf{RIEP}(n-1)$ such that (1) holds.*
- (iv) *There exists a solution B for $\mathbf{RIEP}(n-1)$ such that (1) holds.*

Proof. Note that since (27) holds, we always have (26). We now prove the equivalence between the four statements of the theorem.

(i) \implies (ii) is trivial.

(ii) \implies (iv) by Theorem 2.

(iv) \implies (iii). Since (27) holds, it follows that $\frac{B+B^T}{2}$ also solves $\mathbf{RIEP}(n-1)$.

(iii) \implies (i). Since B is symmetric and since we have (27), the implications (1) and (2) are identical. Our claim now follows by Theorem 30. \square

For uniqueness we have

Theorem 33 *Let $n \geq 2$ and assume that (27) holds. The following are equivalent:*

- (i) *There exists a unique symmetric solution for $\mathbf{RIEP}(n)$.*
- (ii) *There exists a unique solution for $\mathbf{RIEP}(n)$.*
- (iii) *We have $l_{i,i} \neq 0$, $i = 1, \dots, n$.*

Proof. In view of (27), the equivalence of (ii) and (iii) follows from Theorem 4. To see that (i) and (iii) are equivalent note that, by the construction in Theorem 30, for every symmetric solution B for $\mathbf{RIEP}(n-1)$ there exists a solution A for $\mathbf{RIEP}(n)$ such that $A\langle n-1 \rangle = B$. Furthermore, A is uniquely determined if and only if $l_{n,n} \neq 0$. Therefore, it follows that there exists a unique symmetric solution for $\mathbf{RIEP}(n)$ if and only if there exists a unique symmetric solution for $\mathbf{RIEP}(n-1)$ and $l_{n,n} \neq 0$. Our assertion now follows by induction on n . \square

We conclude this section remarking that a similar discussion can be carried over for complex Hermitian matrices.

7 The positive semidefinite case

In view of the discussion of the previous section, it would be interesting to find conditions for the existence of a *positive (semi)definite* real symmetric solution for $\mathbf{RIEP}(n)$. Clearly, a necessary condition is nonnegativity of the numbers s_i whenever $r_i \neq 0$ or $l_i \neq 0$, $i = 1, \dots, n$. Nevertheless, this condition is not sufficient even if a real symmetric solution exists, as is demonstrated by the following example.

Example 34 Let

$$s_1 = 1, \quad s_2 = 3, \quad s_3 = 5,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

The unique solution for **RIEP**(3) is the symmetric matrix

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix},$$

which is not positive semidefinite.

The following necessary and sufficient condition follows immediately from Theorem 4.

Theorem 35 *Let $n \geq 2$ and assume that (27) holds. Assume, further, that $r_{i,i} \neq 0$, $i = 1, \dots, n$. Then the unique solution for **RIEP**(n) is positive semidefinite [positive definite] if and only if $S_n \circ (R_n^T R_n)$ is positive semidefinite [positive definite].*

Remark 36 By Theorem 33, in the case that $r_{i,i} = 0$ for some i we do not have uniqueness of symmetric solutions for **RIEP**(n). Hence, if there exists a symmetric solution for **RIEP**(n) then there exist at least two different such solutions A and B . Note that $A + c(B - A)$ also forms a symmetric solution for **RIEP**(n) for every real number c . It thus follows that in this a case it is impossible to have *all* solutions for **RIEP**(n) positive semidefinite. Therefore, in this case we are looking for conditions for the existence of *some* positive semidefinite solution for **RIEP**(n).

The following necessary condition follows immediately from Proposition 3.

Theorem 37 *Let $n \geq 2$ and assume that (27) holds. If there exists a positive semidefinite real symmetric solution for **RIEP**(n) then $S_n \circ (R_n^T R_n)$ is positive semidefinite.*

In order to find sufficient conditions for the existence of a positive semidefinite solution for **RIEP**(n), we denote by $\sigma(A)$ the least eigenvalue of a real symmetric matrix A .

Lemma 38 *Let $n \geq 2$ and assume that (27) holds. Assume that there exists a symmetric solution A for **RIEP**(n) such that $\sigma(A \langle n-1 \rangle) > s_n$. If $r_n \neq 0$ then $\sigma(A) = s_n$.*

Proof. Since $r_n \neq 0$ it follows that s_n is an eigenvalue of A . Assume that $\sigma(A) \neq s_n$. It follows that A has at least two eigenvalues smaller than or equal to s_n . By the Cauchy Interlacing Theorem for Hermitian matrices, e.g. [8, Theorem 4.3.8, p. 185], it follows that $\sigma(A\langle n-1 \rangle) \leq s_n$, which is a contradiction. Therefore, we have $\sigma(A) = s_n$. \square

Corollary 39 *Let $n \geq 2$ and assume that (27) holds. If $r_i \neq 0$ for all i , $i = 1, \dots, n$, and if $s_1 > s_2 > \dots > s_n \geq 0$ then every real symmetric solution A for **RIEP**(n) is positive semidefinite. If $s_n > 0$ then every real symmetric solution for **RIEP**(n) is positive definite.*

Proof. Note that $A\langle 1 \rangle = [s_1]$ and so $\sigma(A\langle 1 \rangle) = s_1$. Using Lemma 38 repeatedly we finally obtain $\sigma(A) = s_n$, implying our claim. \square

Remark 40 In view of Remark 36, it follows from Corollary 39 that if $r_i \neq 0$ for all i and if $s_1 > s_2 > \dots > s_n \geq 0$ then $r_{i,i} \neq 0$, $i = 1, \dots, n$, and so **RIEP**(n) has a unique (positive semidefinite) solution.

The converse of Corollary 39 is, in general, not true. That is, even if every real symmetric solution for **RIEP**(n) is positive semidefinite we do not necessarily have $s_1 > s_2 > \dots > s_n \geq 0$, as is demonstrated by the following example.

Example 41 Let

$$s_1 = 2, \quad s_2 = 3,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

The unique solution for **RIEP**(3) is the positive definite matrix

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}.$$

Nevertheless, we do not have $s_1 \geq s_2$.

We conclude this section with a conjecture motivated by Theorems 35 and 37. One direction of the conjecture is proven in Theorem 37.

Conjecture 42 *Let $n \geq 2$ and assume that (27) holds. Then there exists a positive semidefinite [positive definite] real symmetric solution for **RIEP**(n) if and only if $S_n \circ (R_n^T R_n)$ is positive semidefinite [positive definite].*

8 The Stieltjes matrix case

In this section we combine the results of the previous two sections to obtain analogous results for *Stieltjes matrices*, that is, symmetric M -matrices.

The following theorem follows immediately from Theorems 30 and 25.

Theorem 43 *Let $n \geq 2$. There exists a symmetric Z -matrix solution for $\mathbf{RIEP}(n)$ if and only if there exists a symmetric Z -matrix solution B for $\mathbf{RIEP}(n-1)$ satisfying*

$$\begin{cases} \frac{s_n \tilde{r}_n}{r_{n,n}} \leq \frac{B \tilde{r}_n}{r_{n,n}}, & r_{n,n} \neq 0 \\ s_n \tilde{r}_n = B \tilde{r}_n, & r_{n,n} = 0 \end{cases},$$

$$\begin{cases} \frac{\tilde{l}_n^T s_n}{l_{n,n}} \leq \frac{\tilde{l}_n^T B}{l_{n,n}}, & l_{n,n} \neq 0 \\ \tilde{l}_n^T s_n = \tilde{l}_n^T B, & l_{n,n} = 0 \end{cases},$$

and

$$l_{n,n} r_{n,n} \neq 0 \implies (s_n I_{n-1} - B) \left(\frac{\tilde{l}_n}{l_{n,n}} - \frac{\tilde{r}_n}{r_{n,n}} \right) = 0.$$

Furthermore, if l_n or r_n is positive then a symmetric Z -matrix solution for $\mathbf{RIEP}(n)$ is a *Stieltjes matrix* if and only if $s_n \geq 0$.

Corollary 44 *Let $n \geq 2$, and assume that the vectors l_i , $i = 1, \dots, n$ are all positive and that (27) holds. There exists a symmetric Z -matrix solution A for $\mathbf{RIEP}(n)$ if and only if there exists a symmetric Z -matrix solution B for $\mathbf{RIEP}(n-1)$ satisfying $s_n \tilde{r}_n \leq B \tilde{r}_n$. The solution A is a *Stieltjes matrix* if and only if $s_n \geq 0$.*

The following nonrecursive sufficient condition from Theorem 27.

Theorem 45 *Let $n \geq 2$, and assume that the vectors l_i , $i = 1, \dots, n$ are all positive, that (27) holds, and that the numbers s_1, \dots, s_n are all positive. If we have*

$$s_i M_i^r \leq s_{i-1} m_i^r, \quad i = 2, \dots, n$$

then there exists a (unique) *Stieltjes matrix* solution A for $\mathbf{RIEP}(n)$.

Proof. By Theorem 27 there exists a unique M -matrix solution A for $\mathbf{RIEP}(n)$. Since A^T also solves the problem, it follows that $A = A^T$ and the result follows.

□

9 The inverse M -matrix case

It is well known that for a nonsingular M -matrix A we have $A^{-1} \geq 0$. Accordingly, a nonnegative matrix A is called *inverse M -matrix* if it is invertible and A^{-1} is an M -matrix. An overview of characterizations of nonnegative matrices that are inverse M -matrices can be found in [9]. In this section we discuss, as a final special case, inverse M -matrix solutions for **RIEP**(n).

The following theorem follows immediately from two results of [9].

Theorem 46 *Let $A \in \mathbb{R}^{n,n}$ be partitioned as in (3). Then A is an inverse M -matrix if and only if B is an inverse M -matrix and*

$$v = B^{-1}y \geq 0, \quad (28)$$

$$u^T = x^T B^{-1} \geq 0, \quad (29)$$

$$s = z - u^T B v > 0, \quad (30)$$

and

$$vu^T \leq -sB^{-1}, \quad \text{except for the diagonal entries.} \quad (31)$$

Proof. By Corollary 3 in [9], if A is an inverse M -matrix then B is an inverse M -matrix. By Theorem 8 in [9], if B is an inverse M -matrix then A is an inverse M -matrix if and only if (28)–(31) hold. Our claim follows. \square

The next result gives necessary and sufficient recursive conditions for the existence of an inverse M -matrix solution for **RIEP**(n).

Theorem 47 *Let $n \geq 2$. There exists an inverse M -matrix solution for **RIEP**(n) if and only if $s_n > 0$ and there exists an inverse M -matrix solution B for **RIEP**($n-1$) satisfying*

$$\begin{cases} \frac{N\tilde{r}_n}{r_{n,n}} \geq 0, & r_{n,n} \neq 0 \\ N\tilde{r}_n = 0, & r_{n,n} = 0 \end{cases}, \quad (32)$$

$$\begin{cases} \frac{\tilde{l}_n^T N}{l_{n,n}} \geq 0, & l_{n,n} \neq 0 \\ \tilde{l}_n^T N = 0, & l_{n,n} = 0 \end{cases}, \quad (33)$$

$$l_{n,n}r_{n,n} \neq 0 \implies \frac{\tilde{l}_n^T N \tilde{r}_n}{l_{n,n}r_{n,n}} < 1, \quad (34)$$

and, except for the diagonal entries,

$$l_{n,n}r_{n,n} \neq 0 \implies s_n \left(\frac{\tilde{l}_n^T N \tilde{r}_n}{l_{n,n}r_{n,n}} - 1 \right) B^{-1} \geq \frac{N\tilde{r}_n \tilde{l}_n^T N}{l_{n,n}r_{n,n}}, \quad (35)$$

where $N = s_n B^{-1} - I_{n-1}$.

Proof. As in the proof of Theorem 2, partition A as in (3). If A is an inverse M -matrix solution for $\mathbf{RIEP}(n)$ then, as is well known, its eigenvalues lie in the open right half plane, and so the real eigenvalue s_n must be positive. Furthermore, by Theorem 46, B is an inverse M -matrix and (28)–(31) hold. Finally, we have (4)–(7). Distinguish between four cases:

1. $r_{n,n} = 0, l_{n,n} \neq 0$. Here x is given by (8), and so it follows from (29) that $\frac{\tilde{l}_n^T N}{l_{n,n}} \geq 0$. By Theorem 2 we have $B\tilde{r}_n = s_n\tilde{r}_n$, implying that $N\tilde{r}_n = 0$.
2. $l_{n,n} = 0, r_{n,n} \neq 0$. Here y is given by (9), and so it follows from (28) that $\frac{N\tilde{r}_n}{r_{n,n}} \geq 0$. By Theorem 2 we have $\tilde{l}_n^T N = 0$.
3. $l_{n,n} = r_{n,n} = 0$. Similarly to the previous cases prove that $N\tilde{r}_n = 0$ and $\tilde{l}_n^T N = 0$.
4. $l_{n,n} \neq 0, r_{n,n} \neq 0$. Here x is given by (8), y is given by (9), and z is given by (10). It follows from (28) that $\frac{N\tilde{r}_n}{r_{n,n}} \geq 0$, and from (29) that $\frac{\tilde{l}_n^T N}{l_{n,n}} \geq 0$. It follows from (30) that

$$\begin{aligned} s &= z - u^T Bv \\ &= s_n - \frac{\tilde{l}_n^T (s_n I_{n-1} - B)\tilde{r}_n}{l_{n,n} r_{n,n}} - \frac{\tilde{l}_n^T (s_n I_{n-1} - B)}{l_{n,n}} B^{-1} B B^{-1} \frac{(s_n I_{n-1} - B)\tilde{r}_n}{r_{n,n}} \\ &= s_n \left(1 - \frac{\tilde{l}_n^T N \tilde{r}_n}{l_{n,n} r_{n,n}} \right) > 0. \end{aligned}$$

Since $s_n > 0$, it now follows that $\frac{\tilde{l}_n^T N \tilde{r}_n}{l_{n,n} r_{n,n}} < 1$. Finally, it follows from (31) that, except for the diagonal entries,

$$\begin{aligned} \frac{N\tilde{r}_n \tilde{l}_n^T N}{l_{n,n} r_{n,n}} &= B^{-1} \frac{(s_n I_{n-1} - B)\tilde{r}_n}{r_{n,n}} \frac{\tilde{l}_n^T (s_n I_{n-1} - B)}{l_{n,n}} B^{-1} = v u^T \\ &\leq -s B^{-1} = s_n \left(\frac{\tilde{l}_n^T N \tilde{r}_n}{l_{n,n} r_{n,n}} - 1 \right) B^{-1}. \end{aligned}$$

We have thus proven that if A is an inverse M -matrix solution for $\mathbf{RIEP}(n)$ then $s_n > 0$ and B is an inverse M -matrix solution B for $\mathbf{RIEP}(n-1)$ satisfying (32)–(35).

Conversely, assume that $s_n > 0$ and B is an inverse M -matrix solution B for $\mathbf{RIEP}(n-1)$ satisfying (32)–(35). We show that x, y and z can be chosen such that (28)–(31) hold, and so by Theorem 46, A is an inverse M -matrix. Here too we distinguish between four cases:

1. $r_{n,n} = 0$, $l_{n,n} \neq 0$. Here x is given by (8), and by (33) we obtain (29). Note that y can be chosen arbitrarily, and z should be chosen such that (7) holds. If we choose $y = 0$ then we obtain (28) and $z = s_n$. It follows that $z - u^T B v = s_n > 0$, and so we also have (30). Finally, since $v = 0$, since $s > 0$ and since B^{-1} is an M -matrix, it follows that (31) holds (except for the diagonal entries).
2. $l_{n,n} = 0$, $r_{n,n} \neq 0$. Here y is given by (9), and by (32) we obtain (28). The vector x can be chosen arbitrarily, so we choose $x = 0$. The proof follows as in the previous case.
3. $l_{n,n} = r_{n,n} = 0$. Here x and y should be chosen such that $x^T \tilde{r}_n = \tilde{l}_n^T y = 0$ and z can be chosen arbitrarily. We choose $x = y = 0$ and the proof follows.
4. $l_{n,n} \neq 0$, $r_{n,n} \neq 0$. Here x is given by (8), y is given by (9), and z is given by (10). By (32) and (33) we obtain (28) and (29) respectively. Finally, similarly to the corresponding case in the proof of the other direction, (34) implies (30) and (35) implies (31). \square

Note that Conditions (32)–(33) imply immediately Conditions (16)–(17) by multiplying the inequality by the nonnegative matrix B . This is not surprising, since an inverse M -matrix is a nonnegative matrix. The converse, however, does not hold in general. The following example shows that although (16)–(17) is satisfied, (32)–(33) do not hold.

Example 48 Let

$$s_1 = 2, \quad s_2 = 5.2361, \quad s_3 = 21.2552,$$

and let

$$l_1 = r_1 = \begin{bmatrix} 1 \end{bmatrix}, \quad l_2 = r_2 = \begin{bmatrix} 0.5257 \\ 0.8507 \end{bmatrix}, \quad l_3 = r_3 = \begin{bmatrix} 0.1349 \\ 0.3859 \\ 0.9126 \end{bmatrix}.$$

The unique solution for **RIEP**(3) is the nonnegative matrix

$$A = \begin{bmatrix} 2 & 2 & 2 \\ 2 & 4 & 7 \\ 2 & 7 & 18 \end{bmatrix}$$

which is not an inverse M -matrix since

$$A^{-1} = \begin{bmatrix} 1.6429 & -1.5714 & 0.4286 \\ -1.5714 & 2.2857 & -0.7143 \\ 0.4286 & -0.7143 & 0.2857 \end{bmatrix}.$$

Indeed, the unique nonnegative solution $B = \begin{bmatrix} 2 & 2 \\ 2 & 4 \end{bmatrix}$ for **RIEP**(2) satisfies (16), as

$$s_3 \tilde{r}_3 = \begin{bmatrix} 2.8673 \\ 8.2024 \end{bmatrix} \geq \begin{bmatrix} 1.0416 \\ 1.8134 \end{bmatrix} = B \tilde{r}_3.$$

However, B does not satisfy (32), since the vector

$$N \tilde{r}_3 = (s_3 B^{-1} - I_2) \tilde{r}_3 = \begin{bmatrix} -1.3688 \\ 2.2816 \end{bmatrix}$$

is not nonnegative.

References

- [1] A. Ben-Israel and T.N.E. Greville, *Generalized Matrix Inverses: Theory and Applications*, John Wiley, New York, 1974.
- [2] A. Berman and R. Plemmons, *Nonnegative Matrices in Mathematical Sciences*, SIAM, Philadelphia, 1994.
- [3] D. Boley and G.H. Golub, A survey of matrix inverse eigenvalue problems, *Inverse problems* **3** (1987), 595–622.
- [4] M. Boyle and D. Handelmann, The spectra of non-negative matrices via symbolic dynamics, *Ann. Math.* **133** (1991), 249–316.
- [5] S. Friedland, On an inverse problem for nonnegative and eventually nonnegative matrices, *Israel J. Math.* **29** (1978), 43–60.
- [6] G. Frobenius, Über Matrizen aus positiven Elementen, *S.-B. Preuss. Akad. Wiss.* (1909), 471–476.
- [7] H. Hochstadt, On some inverse problems in matrix theory, *Arch. Math.* **18** (1967), 201–207.
- [8] R.A. Horn and C.R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- [9] C.R. Johnson, Inverse M -matrices, *Lin. Alg. Appl.* **47** (1982), 195–216.
- [10] D.M. Koteljanskiĭ, On some properties of matrices with positive elements, *Transl. Amer. Math. Soc. Ser. 2* **27** (1963), 9–18.
- [11] T.J. Laffey, Inverse eigenvalue problems for matrices, *Proc. Royal Irish Acad.* **95A** (suppl.) (1995), 81–88.

- [12] R. Loewy and D. London, A note on an inverse problem for nonnegative matrices, *Lin. and Multilin. Alg.* **6** (1978), 83–90.
- [13] T. Markham, Nonnegative matrices whose inverses are M -matrices, *Proc. Amer. Math. Soc.* **36** (1972), 326–330.
- [14] G.N. de Oliveira, Note on an inverse characteristic value problem, *Numer. Math.* **15** (1970), 345–347.

Other titles in the SFB393 series:

- 99-01 P. Kunkel, V. Mehrmann, W. Rath. Analysis and numerical solution of control problems in descriptor form. January 1999.
- 99-02 A. Meyer. Hierarchical preconditioners for higher order elements and applications in computational mechanics. January 1999.
- 99-03 T. Apel. Anisotropic finite elements: local estimates and applications (Habilitationsschrift). January 1999.
- 99-04 C. Villagonzalo, R. A. Römer, M. Schreiber. Thermoelectric transport properties in disordered systems near the Anderson transition. February 1999.
- 99-05 D. Michael. Notizen zu einer geometrisch motivierten Plastizitätstheorie. Februar 1999.
- 99-06 T. Apel, U. Reichel. SPC-PM Po 3D V 3.3, User's Manual. February 1999.
- 99-07 F. Tröltzsch, A. Unger. Fast solution of optimal control problems in the selective cooling of steel. March 1999.
- 99-08 W. Rehm, T. Unger (Eds.). Ausgewählte Beiträge zum 2. Workshop Cluster-Computing 25./26. März 1999, Universität Karlsruhe. März 1999.

The complete list of current and former preprints is available via
<http://www.tu-chemnitz.de/sfb393/preprints.html>.