

IRREGULAR HODGE FILTRATION OF SOME CONFLUENT HYPERGEOMETRIC SYSTEMS

ALBERTO CASTAÑO DOMÍNGUEZ AND CHRISTIAN SEVENHECK

ABSTRACT. We calculate the irregular Hodge filtration, recently introduced in [Sab15], for the purely irregular hypergeometric \mathcal{D} -modules.

1. INTRODUCTION

The aim of this paper is to compute some Hodge theoretic invariants of certain classical differential systems in one variable. These are the so-called irregular Hodge numbers, which have been introduced recently by Sabbah [Sab15]. They are called irregular because they are attached to differential systems which may have irregular singular points, a feature that is excluded for classical variation of Hodge structures as well as for the more general Hodge modules. The very definition of these numbers rely on the theory of mixed twistor modules of T. Mochizuki ([Moc15a]). Twistor modules generalise Hodge modules, in the sense that the underlying \mathcal{D} -module of a twistor module can have irregular singularities. In particular, one can define a version of the Fourier-Laplace transformation functor within the category of mixed twistor modules, which is impossible for mixed Hodge modules. The drawback of this generalization is that one cannot directly assign a filtration to a twistor module, and hence it is not easy to attach numerical invariants, like Hodge numbers to it. In the above mentioned paper [Sab15], Sabbah constructs an intermediate category between mixed Hodge modules and mixed twistor modules (called irregular mixed Hodge modules) which is on the one hand sufficiently large to be stable under all relevant operations that are defined for twistor modules (in particular, the Fourier-Laplace transformation), but which allows one to define a filtration, called irregular Hodge filtration, for each of its objects. The construction is related to the earlier papers [Yu14, ESY17, SY15], where for certain projective morphisms $f : X \rightarrow \mathbb{P}^1$, a (rationally indexed) filtration was introduced on the twisted de Rham complex $(\Omega^\bullet(*D), d + df \wedge)$, where $D \subset X$ is a certain normal crossing boundary divisor such that $f|_{X \setminus D}$ yields a regular function. Instead of considering meromorphic differential forms, one can also use the subcomplex of so-called f -adapted logarithmic forms (also called Kontsevich complex, see [KKP17]) Ω_f^\bullet , which consists of forms such that the exterior product with df is still logarithmic along D . From an E_1 -degeneration property of the corresponding spectral sequence proved in [ESY17], one obtains a filtration on the twisted de Rham cohomology, called irregular Hodge filtration of the regular function $f : X \setminus D \rightarrow \mathbb{A}^1$. We refer to [ESY17] for more details.

Notice also that the recent paper [KKP17] gives three definitions of so-called Landau-Ginzburg Hodge numbers associated to a family $f : X \rightarrow \mathbb{P}^1$, one of them being $\dim H^p(\Omega_f^q)$. Conjecturally these three definitions coincide, but this seems to require some more assumptions (see [Sha17] for some partial results). Although these Hodge numbers have only integer indices, they are closely related to the dimensions of the graded parts of the filtration form [ESY17]. Ultimately, according to [KKP17] and following prediction from homological mirror symmetry, one hopes for a correspondence between the Hodge numbers of some, say, projective varieties entering in the A-model and the irregular Hodge numbers of its (Landau-Ginzburg) B-model.

For the moment, there are quite a few examples where the irregular Hodge numbers (in any of its above mentioned variants) can actually be computed. A central result of [Sab15] states that

2010 *Mathematics Subject Classification*. Primary 14F10, 32C38.

The authors are partially supported by the project SISYPH: ANR-13-IS01-0001-01/02 and DFG grant SE 1114/5-1.

rigid irreducible \mathcal{D} -modules on the projective line underly objects of the category of irregular mixed Hodge modules, and consequently admit a unique irregular Hodge filtration, provided that their formal local monodromies are unitary. Rigid \mathcal{D} -modules are particularly interesting since they can be algorithmically constructed from simple objects by an algorithm due to Arinkin and Katz (cf. [Ari10]). Among the most studied and best understood examples of such rigid \mathcal{D} -modules are the classical hypergeometric \mathcal{D} -modules. In the regular case, Fedorov has recently given in [Fed15] a closed formula for the Hodge numbers using the work [DS13] of Dettweiler and Sabbah.

The present papers aims at extending Fedorov's work to some special class of irregular hypergeometric modules, where we can explicitly control the (irregular) Hodge filtration and calculate the (irregular) Hodge numbers. Our principal result, Theorem 4.7, gives a very simple formula for these numbers, which is in some sense similar to the shape of Fedorov's formula. The main ingredients are a reduction process (as explained in [BZMW13]) to obtain classical hypergeometric \mathcal{D} -modules from some higher dimensional ones, the so-called GKZ-systems, techniques from [Rei14] and [RS15b, RS17] (going back to [Sab08]) to understand Hodge module structures on these GKZ-systems as well as a quite explicit solution to the so-called Birkhoff problem that is inspired by calculations in toric mirror symmetry (see again [RS15b] as well as [DS04] and also [GMS09]).

While writing this paper, we learned that the same formula has been proved by Sabbah and Yu in the newest version of [Sab15] by different means. Nevertheless, we believe that the geometric approach taken here has some potential to be generalized to arbitrary irregular hypergeometric systems (see Conjecture 3.23 below).

Let us recall some notation that will be adopted throughout the paper: For a smooth complex algebraic variety X , we write \mathcal{D}_X for the sheaf of algebraic differential operators on X . If X is affine, we sometimes switch freely between sheaves of \mathcal{D}_X -modules and modules of global sections. For a morphism $f : X \rightarrow Y$, we denote the direct resp. inverse image functors for \mathcal{D} -modules as usual by f_+ resp. f^+ (see [HTT08] for a thorough discussion of these and related functors). We put $\mathbb{G}_m := \text{Spec } \mathbb{C}[t, t^{-1}]$; if we want to fix a coordinate on this one-dimensional torus, we also write $\mathbb{G}_{m,t}$. We denote by $\mathcal{R}_{\mathbb{A}_z^1 \times X}^{\text{int}}$ the sheaf of Rees rings on X (with integrable structure), that is, the subsheaf of non-commutative algebras of $\mathcal{D}_{\mathbb{A}_z^1 \times X}$ locally generated by $z^2 \partial_z$ and $(z \partial_{x_i})_{i=1, \dots, n}$, where (x_1, \dots, x_n) are local coordinates on X . We let $\text{MHM}(X)$ resp. $\text{MTM}(X)$ be the abelian categories of algebraic mixed Hodge modules (see [Sai90]) resp. algebraic mixed twistor modules (see [Moc15a]) on X . The category $\text{MTM}^{\text{int}}(X)$ consists of those mixed twistor modules where the underlying $\mathcal{R}_{\mathbb{A}_z^1 \times X}$ -modules have an integrable structure, i.e., where they are modules over $\mathcal{R}_{\mathbb{A}_z^1 \times X}^{\text{int}}$. The category $\text{IrrMHM}(X)$, as defined in [Sab15, Def. 2.52], is a certain subcategory of $\text{MTM}^{\text{int}}(X)$ (or rather of a variant, called ${}_{\iota}\text{MTM}^{\text{int}}(X)$, which is shown to be equivalent to $\text{MTM}^{\text{int}}(X)$ in op. cit.) consisting of objects that satisfy a so-called rescaling condition and a grading property. Its main feature is that the \mathcal{D}_X -module \mathcal{M} associated to an object $\widehat{\mathcal{M}}$ in $\text{IrrMHM}(X)$ carries a good filtration $F_{\bullet} \mathcal{M}$, indexed by \mathbb{R} , called the irregular Hodge filtration.

Acknowledgements. We would like to thank Takuro Mochizuki, Thomas Reichelt and Claude Sabbah for their interest in our work and for many stimulating discussions. We thank the Max Planck Institute for Mathematics in the Sciences, where a significant part of the work presented here has been carried out.

2. HYPERGEOMETRIC \mathcal{D} -MODULES AND DIMENSIONAL REDUCTIONS

In this section we will introduce two different kinds of hypergeometric \mathcal{D} -modules: classical and GKZ. We will dedicate more time to classical ones, since they form one of the main objects of study of the paper and will end by showing the relation between both types, which will be useful for us in the future.

Definition 2.1. Let $(n, m) \neq (0, 0)$ be a pair of nonnegative integers, and let $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_m be elements of \mathbb{C} . The (classical) hypergeometric \mathcal{D} -module associated with the α_i and the β_j is defined

as the quotient of $\mathcal{D}_{\mathbb{G}_m}$ by the ideal generated by the so called hypergeometric operator

$$\prod_{i=1}^n (t\partial_t - \alpha_i) - t \prod_{j=1}^m (t\partial_t - \beta_j).$$

We will denote it by $\mathcal{H}(\alpha_1, \dots, \alpha_n; \beta_1, \dots, \beta_m)$, or in an abridged way, $\mathcal{H}(\alpha_i; \beta_j)$.

Remark 2.2. The excluded type $(n, m) = (0, 0)$ corresponds to the punctual delta $\mathcal{D}_{\mathbb{G}_m}$ -module on \mathbb{G}_m $\mathcal{H}(\emptyset; \emptyset) = \mathcal{D}_{\mathbb{G}_m}/(1-x)$.

On the other hand, if we denote the Kummer \mathcal{D} -module $\mathcal{D}_{\mathbb{G}_m}/(t\partial_t - \eta)$ by \mathcal{K}_η , for any fixed complex number η , then $\mathcal{H}(\alpha_i; \beta_j) \otimes_{\mathcal{O}_{\mathbb{G}_m}} \mathcal{K}_\eta \cong \mathcal{H}(\alpha_i + \eta; \beta_j + \eta)$.

Every hypergeometric \mathcal{D} -module has Euler characteristic -1 (cf. [Kat90, Lem. 2.9.13]). For $n \neq m$, no hypergeometric \mathcal{D} -module of type (n, m) has singularities on \mathbb{G}_m . If $n > m$ (resp. $m > n$), they have a regular singularity at the origin (resp. infinity) and an irregular singularity at infinity (resp. the origin) of irregularity one and slope $1/|n-m|$ of multiplicity $|n-m|$ (cf. [Kat90, Prop. 2.11.9]). For $n = m$, the hypergeometric modules of type (n, n) are regular, with singularities only at the origin, infinity and 1.

Proposition 2.3 (Irreducibility). (cf. [Kat90, Prop. 2.11.9, 3.2]) *Let $\mathcal{H} := \mathcal{H}(\alpha_i; \beta_j)$ be a hypergeometric \mathcal{D} -module. It is irreducible if and only if for any pair (i, j) of indices, $\alpha_i - \beta_j$ is not an integer.*

Proposition 2.4 (Rigidity). *Let $\mathcal{H} := \mathcal{H}(\alpha_i; \beta_j)$ be an irreducible hypergeometric \mathcal{D} -module of type (n, m) , where $n \geq m$, and let \mathcal{M} be another irreducible $\mathcal{D}_{\mathbb{G}_m}$ -module of Euler-Poincaré characteristic -1 which has no singularities outside $\{0, 1, \infty\}$. Assume that*

- $\mathcal{H} \otimes \mathbb{C}((t)) \cong \mathcal{M} \otimes \mathbb{C}((t))$.
- $\mathcal{H} \otimes \mathbb{C}((1/t)) \cong \mathcal{M} \otimes \mathbb{C}((1/t))$.
- If $n = m$, assume further that \mathcal{M} has a regular singularity at 1.

In that case, \mathcal{H} and \mathcal{M} are isomorphic.

Proof. \mathcal{H} and \mathcal{M} being irreducible, both of them coincide with the middle extension of their restriction to $\mathbb{G}_m \setminus \{1\}$. Assume first that $n = m$, that is, \mathcal{H} is regular. Since \mathcal{M} has characteristic -1 and regular singularities at the origin, one and infinity, by the formula for the Euler characteristic [Kat90, Thm. 2.9.9], its formal local monodromy at 1 must be a pseudoreflection. Then we can apply [op. cit., Rigidity Thm. 3.5.4] and we are done.

If $n > m$, \mathcal{M} , like \mathcal{H} , has a regular singularity at the origin and an irregular singularity at infinity. By the same formula for the Euler characteristic of $j_+ j^+ \mathcal{M}$ (denoting by j the inclusion $\mathbb{G}_m \setminus \{1\} \hookrightarrow \mathbb{G}_m$), \mathcal{M} cannot have more singularities in \mathbb{P}^1 apart from zero and infinity. In that case we apply [op. cit., Rigidity Thm. bis 3.7.3]. \square

Consider the case in which $\mathcal{H} = \mathcal{H}(\alpha_i; \beta_j)$ is of type (n, n) , i.e., such that it is regular. From the rigidity property of the last Proposition, one concludes by [Sim90, Cor. 8.1] that the restriction of \mathcal{H} to $\mathbb{G}_m \setminus \{1\}$ underlies a complex polarizable variation of Hodge structures. The next statement gives important information on its Hodge invariants.

Proposition 2.5 (Hodge numbers (regular case)). (cf. [Fed15, Thm. 1]) *Let $\mathcal{H} = \mathcal{H}(\alpha_i; \beta_j)$ be an irreducible hypergeometric \mathcal{D} -module of type (n, n) . Assume that the α_i and the β_j are increasingly ordered real numbers, lying in the interval $[0, 1)$. Set*

$$\rho(k) = |\{j = 1, \dots, n : \alpha_j < \beta_k\}| - k,$$

for $k = 1, \dots, n$. Then the Hodge numbers of \mathcal{H} are, up to a shift,

$$h^p = |\rho^{-1}(p)| = |\{k = 1, \dots, n : \rho(k) = p\}|.$$

This last result is the analogous one, in the regular case, to theorem 4.7, and served as the main motivation to start this project. Now that we have seen part of the behaviour of classical hypergeometric \mathcal{D} -modules, let us continue with the other family of hypergeometric modules mentioned above.

Definition 2.6. Let $n \geq m$ two positive integers, and let $d = n - m$. Let $\beta \in \mathbb{C}^d$ be a vector and let $A \in M(d \times n, \mathbb{Z})$ be an integer matrix. Consider the affine space \mathbb{A}^n with coordinates $\lambda_1, \dots, \lambda_n$. We define the Euler operators $E_i = \sum_j a_{ij} \lambda_j \partial_{\lambda_j}$, for $i = 1, \dots, d$, and the toric ideal

$$I_A := (\partial_\lambda^u - \partial_\lambda^v : Au = Av) \subset \mathbb{C}[\partial_{\lambda_1}, \dots, \partial_{\lambda_n}] \subset D_{\mathbb{A}^n}.$$

Then, the GKZ-hypergeometric \mathcal{D} -module is $\mathcal{M}_A^\beta := \mathcal{D}_{\mathbb{A}^n} / (\mathcal{I}_A + (E_i - \beta_i : i = 1, \dots, d)_{i=1, \dots, d})$.

We call homogenization of a GKZ-system \mathcal{M}_A^β the GKZ-hypergeometric system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$, where \tilde{A} is the extended matrix

$$\left(\begin{array}{c|c} 1 & \underline{1}_n^t \\ \hline \underline{0}_d & A \end{array} \right)$$

and $\tilde{\beta} = (\beta_0, \beta_1, \dots, \beta_d)^t$, for certain $\beta_0 \in \mathbb{C}$ (here and later in this paper, we write \underline{x}_k for a column vector with k rows, where all entries contain the value x).

A classical hypergeometric module can be considered as a dimensional reduction of a certain GKZ-system. We will describe this procedure in some more detail now, because it allows us to apply some of the many known results on GKZ-systems to classical hypergeometric modules.

Proposition 2.7. *Let m be a positive integer. Let $A \in M((m-1) \times m, \mathbb{Z})$ be an integer matrix of rank $m-1$ and take $\kappa \in \mathbb{C}^m$ such that $\kappa_1 = 0$. Consider the inclusion $\iota : \mathbb{G}_m \hookrightarrow \mathbb{A}^m$ given by $t \mapsto (t, 1, \dots, 1)$, and let $B = (b_1, \dots, b_m) \in \mathbb{Z}^m$ be a Gale dual of A , that is, an integer column matrix which generates $\ker_{\mathbb{Q}} A$. Put*

$$\eta := \prod_{b_i > 0} b_i^{b_i} \prod_{b_j < 0} (-b_j)^{-b_j}.$$

Then we have $h_\eta^+ \mathcal{H}(\alpha_i; \beta_j) \cong \iota^+ \mathcal{M}_A^{\kappa}$, where h_η is the automorphism of \mathbb{G}_m given by $t \mapsto \eta t$, and the unordered sets of parameters α_i and β_j , counted with multiplicities, are

$$(\alpha_i) = \left(\frac{k - \kappa_j}{b_j} : b_j > 0, k = 0, \dots, b_j - 1 \right),$$

$$(\beta_i) = \left(\frac{\kappa_j - k}{b_j} : b_j < 0, k = 0, \dots, b_j - 1 \right).$$

Proof. On one hand, \mathcal{M}_A^{κ} is not only a GKZ-hypergeometric module, but also a lattice basis binomial $\mathcal{D}_{\mathbb{A}^m}$ -module (cf. [BZMW13, 0.2]). This is because A being of codimension one implies that the toric ideal \mathcal{I}_A is a complete intersection, and then coincides with the lattice basis ideal

$$\mathcal{I}(B) := (\partial_\lambda^{w^+} - \partial_\lambda^{w^-} : w = w_+ - w_- \text{ is a column of } B).$$

On the other, the expression we have given for the parameters of $\mathcal{H}(\alpha_i; \beta_j)$ follows from applying the definition of Horn hypergeometric \mathcal{D} -modules given in [op. cit., 0.3] for a column matrix (restricted to the torus $\mathbb{G}_{m,t}$) and compare it with Definition 2.1.

Now let $j : \mathbb{G}_m \hookrightarrow \mathbb{A}^1$ be the canonical inclusion. The isomorphism given in [op. cit., Thm. 8.1] relates lattice basis binomial \mathcal{D} -modules to Horn hypergeometric ones. However, due to the previous discussions, we obtain the isomorphism in the statement just by applying j^+ to both sides of it. \square

In our study of Hodge theoretic properties of hypergeometric $\mathcal{D}_{\mathbb{G}_m}$ -modules below, we need to go in some sense in the opposite direction: given sets $\{\alpha_i\}$, $\{\beta_j\}$, we would like to determine a matrix A and a parameter vector β such that the module $\mathcal{H}(\alpha_i; \beta_j)$ can be obtained as an inverse image of the GKZ-system \mathcal{M}_A^β . Although such an pair (A, β) is not unique, a systematic way of constructing it can be formulated as follows.

Corollary 2.8. *Let $\mathcal{H}(\alpha_i; \beta_j)$ be a hypergeometric $\mathcal{D}_{\mathbb{G}_m}$ -module of type (n, m) with $\alpha_1 = 0$, let $A \in M((m+n-1) \times (n+m), \mathbb{Z})$ given by*

$$A = \left(\begin{array}{c|c|c} \mathbb{1}_m & \mathbb{0}_{m \times (n-1)} & \text{Id}_m \\ \hline \mathbb{1}_{n-1} & -\text{Id}_{n-1} & \mathbb{0}_{(n-1) \times m} \end{array} \right),$$

and let $\beta = (\beta_1, \dots, \beta_m, \alpha_2, \dots, \alpha_n)^t$. Let $\iota: \mathbb{G}_m \rightarrow \mathbb{A}^{n+m}$ by given by $t \mapsto (t, 1, \dots, 1)$. Then

$$\mathcal{H}(\alpha_i; \beta_j) \cong \iota^+ \mathcal{M}_A^\beta.$$

Proof. The statement is an easy consequence of the Proposition, taking $B = (1, \dots, 1, -1, \dots, -1)^t$ and $\kappa = (0, -\alpha_2, \dots, -\alpha_n, \beta_1, \dots, \beta_m)^t$. \square

We will see later that the restriction $\alpha_1 = 0$ is not as strong as it may appear: By tensoring a given hypergeometric module with an appropriate Kummer module, we can always reach this assumption, and often the statements we are after can be shown for Kummer modules in a rather elementary way.

3. HODGE MODULES AND FOURIER-LAPLACE TRANSFORMATION

Let $\alpha_1, \dots, \alpha_n$ be real numbers in $[0, 1)$, and consider the hypergeometric $\mathcal{D}_{\mathbb{G}_m}$ -module $\mathcal{H} = \mathcal{H}(\alpha_i; \emptyset)$. The goal of this section is proving Theorem 3.22 below, showing that \mathcal{H} underlies an object of $\text{IrrMHM}(\mathbb{G}_m)$ (the abelian category of exponential Hodge modules as defined in [Sab15]) that can be extended in a unique way to the projective line, allowing us to compute the irregular Hodge filtration of \mathcal{H} . To reach the aforementioned theorem we will work with some twistor GKZ-hypergeometric systems, following mainly [Moc15b, RS15b, RS17]. Although we will give a direct definition, a comprehensive treatment of that situation can be found in those references.

Let $d < n$ be two positive integers, and take a parameter vector $\beta \in \mathbb{C}^d$. For the Hodge theoretic questions we are interested in, only real parameter vectors are relevant, but we will work with this more general setting until Remark 3.15 below. Let $A = (\underline{a}_1, \dots, \underline{a}_n) \in M(d \times n, \mathbb{Z})$ an integer matrix satisfying the following:

Assumptions 3.1.

- (1) $\mathbb{Z}A = \mathbb{Z}^d$, here $\mathbb{Z}A := \sum_{i=1}^n \mathbb{Z}\underline{a}_i$,
- (2) A is saturated, that is, we have $\mathbb{N}A = (\mathbb{R}_{\geq 0}A) \cap \mathbb{Z}^d$,
- (3) If we denote by $\text{Conv}(\underline{a}_1, \dots, \underline{a}_n)$ the convex hull in \mathbb{R}^d of the vectors given by the columns of the matrix A , then the origin lies strictly in the interior of $\text{Conv}(\underline{a}_1, \dots, \underline{a}_n)$.

As before, we write

$$\left(\begin{array}{c|c} 1 & \mathbb{1}_n^t \\ \hline \mathbb{0}_d & A \end{array} \right)$$

for the extended matrix $\tilde{A} \in M((d+1) \times (n+1), \mathbb{Z})$, and we consider the extended parameter $\tilde{\beta} = (\beta_0, \beta)$ for some $\beta_0 \in \mathbb{C}$. From these data, we could consider the A-hypergeometric systems \mathcal{M}_A^β or $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$, but we will focus on their twistor versions, as commented at the beginning of the section.

Our main case of interest will be given by the matrix

$$(1) \quad A = \begin{pmatrix} 1 & -1 & 0 & 0 & \dots & 0 \\ 1 & 0 & -1 & 0 & \dots & 0 \\ \vdots & \dots & & & & \\ 1 & 0 & 0 & 0 & \dots & -1 \end{pmatrix} \in M((n-1) \times n, \mathbb{Z}),$$

and where the parameter vector is $\beta = \alpha := (\alpha_2, \dots, \alpha_n)^t \in \mathbb{R}^{n-1}$ (these will be the numbers α_i of our given classical hypergeometric system, notice that α_1 is not appearing here). Notice that the above assumptions 3.1 are satisfied in this example.

Let $S_1 = \mathbb{G}_m^d = \text{Spec}(\mathbb{C}[y_1^\pm, \dots, y_d^\pm])$ and $S_2 = \mathbb{G}_m^n = \text{Spec}(\mathbb{C}[\lambda_1^\pm, \dots, \lambda_n^\pm])$ two algebraic tori, and consider the affine space $V = \mathbb{A}^{n+1}$ with coordinates $\lambda_0, \lambda_1, \dots, \lambda_n$. We let V^\vee be the dual space with

coordinates w_0, w_1, \dots, w_n , and we also set $\tau := -w_0$ and $z := \tau^{-1}$. We decompose $V = \mathbb{A}_{\lambda_0}^1 \times W$, and consider $S_2 \subset W$ as an open subset.

Recall that for a smooth algebraic variety X , we denote by $\mathcal{R}_{\mathbb{A}_z^1 \times X}^{\text{int}}$ the sheaf of Rees rings on X with integrable structure. Then we put $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)} = \mathcal{R}_{\mathbb{A}_z^1 \times S_2}^{\text{int}} / \mathcal{I}$, where $\beta_0 \in \mathbb{C}$ and \mathcal{I} is generated by

$$\begin{aligned} & \prod_{j:l_j > 0} (z\partial_{\lambda_j})^{l_j} - \prod_{j:l_j < 0} (z\partial_{\lambda_j})^{-l_j}, \quad \underline{l} \in \mathbb{L} \\ & z^2\partial_z + \lambda_1 z\partial_{\lambda_1} + \dots + \lambda_n z\partial_{\lambda_n} - z\beta_0, \\ & \sum_{j=1}^n a_{kj} \lambda_j z\partial_{\lambda_j} - z\beta_k, \quad k = 1, \dots, d \end{aligned}$$

where $\underline{l} = (l_1, \dots, l_n)$ and $\mathbb{L} \subset \mathbb{Z}^n$ is the kernel of the linear map $\mathbb{Z}^n \rightarrow \mathbb{Z}^d$ given by left multiplication by the matrix A .

In the special case of our original matrix from (1) the generators of \mathcal{I} are

$$\begin{aligned} & (z\partial_{\lambda_1}) \cdot \dots \cdot (z\partial_{\lambda_n}) - 1 \\ & z^2\partial_z + \lambda_1 z\partial_{\lambda_1} + \dots + \lambda_n z\partial_{\lambda_n} - z\beta_0. \\ & \lambda_1 z\partial_{\lambda_1} - \lambda_i z\partial_{\lambda_i} - z\beta_i, \quad i = 2, \dots, n \end{aligned}$$

The following is a variant of [RS15b, Lem. 2.13].

Lemma 3.2. $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ is locally $\mathcal{O}_{\mathbb{A}_z^1 \times S_2}$ -free of rank $n! \text{vol}(\Delta(\underline{a}_1, \dots, \underline{a}_n))$ (considering the normalized volume in \mathbb{R}^n such that $[0, 1]^n$ has volume one). For the case of the matrix in equation (1), this rank equals n .

For any complex number β , recall that the Kummer \mathcal{D} -module of parameter β is by definition the quotient $\mathcal{K}_\beta := \mathcal{D}_{\mathbb{G}_{m,t}} / (t\partial_t - \beta)$. We will also use in this section \mathcal{R} -modules arising from such \mathcal{D} -modules. Here is the precise definition.

Definition 3.3. For any complex number β , we define the Kummer \mathcal{R} -module of parameter β as the cyclic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -module

$$\widehat{\mathcal{K}}_\beta := \mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_{m,t}}^{\text{int}} / (z^2\partial_z, zt\partial_t - z\beta).$$

Remark 3.4. Although both kinds of Kummer modules can be defined for any complex value of their parameters, in the end we will be interested only in the real case to make use of their Hodge properties. Indeed, since both have no singularities at $\mathbb{G}_{m,t}$, if \mathcal{K}_β were a complex Hodge module it would be in fact a complex variation of Hodge structures, and by (the first part of) the proof of [Sch73, Lem. 4.5], β must be real.

On the other hand, $\widehat{\mathcal{K}}_\beta$ is clearly the Rees module of \mathcal{K}_β together with the trivial filtration F_\bullet such that $F_k = 0$ for $k < 0$ and $F_k = \mathcal{K}_\beta$ for $k \geq 0$. As described in [Moc15a, Prop. 13.5.4], it gives rise to an integrable pure twistor $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module, which belongs as well to $\text{IrrMHM}(\mathbb{G}_m)$. It is also described as a harmonic bundle at [op. cit., §2.1.9].

Definition 3.5. For any smooth complex algebraic variety X , and for any $\beta \in \mathbb{C}^d$, we will denote by $\mathcal{O}_{S_1 \times X}^\beta$ the $\mathcal{D}_{S_1 \times X}$ -module corresponding to the structure sheaf of $S_1 \times X$ twisted by $\underline{y}^{-1-\beta}$, that is,

$$\mathcal{O}_{S_1 \times X}^\beta := \frac{\mathcal{D}_{S_1 \times X}}{(y_k \partial_{y_k} + \beta_k + 1 : k = 1, \dots, d)} \boxtimes \mathcal{O}_X =: \mathcal{O}_{S_1}^\beta \boxtimes \mathcal{O}_X.$$

Note that for any other $\beta' \in \mathbb{C}^d$ such that $\beta - \beta' \in \mathbb{Z}^d$, $\mathcal{O}_{S_1 \times X}^\beta \cong \mathcal{O}_{S_1 \times X}^{\beta'}$. These modules underly complex Hodge modules iff the components of the parameter vector are real numbers, since they are the corresponding exterior product of the Kummer modules $\mathcal{K}_{\beta_1}, \dots, \mathcal{K}_{\beta_d}$ and \mathcal{O}_X .

We need a slight variant of a comparison result from [RS15b, Prop. 2.14] and [RS17, Prop. 3.21]. A more general version, which is not needed here, can be found in [Moc15b, Prop. E.4, E.6]. Consider

the following family of Laurent polynomials

$$(2) \quad \begin{aligned} \varphi : S_1 \times S_2 &\longrightarrow \mathbb{A}_{\lambda_0}^1 \times S_2 \\ (\underline{y}, \underline{\lambda}) &\longmapsto \left(-\sum_{i=1}^n \lambda_i \underline{y}^{a_i}, \lambda_1, \dots, \lambda_n \right), \end{aligned}$$

which in the case of the matrix from equation (1) becomes

$$(3) \quad \varphi(y_1, \dots, y_{n-1}, \lambda_1, \dots, \lambda_n) = \left(\frac{\lambda_1}{y_1} + \dots + \frac{\lambda_{n-1}}{y_{n-1}} + \lambda_n \cdot y_1 \cdots y_{n-1}, \lambda_1, \dots, \lambda_n \right).$$

Write $f : S_1 \times S_2 \rightarrow \mathbb{A}_{\lambda_0}^1$ for the composition of φ with the first projection $\mathbb{A}_{\lambda_0}^1 \times S_2 \rightarrow \mathbb{A}_{\lambda_0}^1$. With these notations, the following is proved precisely as in [RS17, Prop. 3.21], taking into account the twist of $\mathcal{O}_{S_1 \times S_2}$ by $y^{-1-\beta}$.

Proposition 3.6. *There exists an isomorphism of $\mathcal{R}_{\mathbb{A}^1 \times S_2}^{\text{int}}$ -modules*

$$\mathcal{H}^0 \left(\pi_{2,*} \Omega_{S_1 \times S_2 / S_2}^{\bullet+d} [z], z(d - \kappa(\beta)\wedge) - df \wedge \right) \longrightarrow \widehat{\mathcal{M}}_A^{(0,\beta)}$$

for any $\beta \in \mathbb{C}^d$, given by sending $\omega := \prod_{j=1}^d y_j / dy_j$ to $[1] \in \widehat{\mathcal{M}}_A^{(0,\beta)}$, where we write $\kappa(\beta)$ for $\sum_{j=1}^d \beta_j dy_j / y_j$ and π_2 for the second canonical projection $S_1 \times S_2 \rightarrow S_2$.

In order to achieve the comparison results of this section we need to use several variants of the Fourier-Laplace transformations; let us recall here the definitions.

Definition 3.7. Let Y be a smooth algebraic variety, U be a finite-dimensional complex vector space and U' its dual vector space. Denote by \mathcal{E} the trivial vector bundle $\tau : U \times Y \rightarrow Y$ and by \mathcal{E}' its dual. Write $\text{can} : U \times U' \rightarrow \mathbb{A}^1$ for the canonical morphism defined by $\text{can}(a, \varphi) = \varphi(a)$. This extends to a function $\text{can} : \mathcal{E} \times \mathcal{E}' \rightarrow \mathbb{A}^1$. Define $\mathcal{L} := \mathcal{O}_{\mathcal{E} \times_Y \mathcal{E}'} e^{-\text{can}}$, the free rank one module with differential given by the product rule. Consider also the canonical projections $p_1 : \mathcal{E} \times_Y \mathcal{E}' \rightarrow \mathcal{E}$, $p_2 : \mathcal{E} \times_Y \mathcal{E}' \rightarrow \mathcal{E}'$. The partial Fourier-Laplace transformation is then defined by

$$\text{FL}_Y := p_{2,+} (p_1^+ \bullet \otimes^{\mathbf{L}} \mathcal{L}).$$

If the base Y is a point we recover the usual Fourier-Laplace transformation and we will simply write FL. Notice that although this functor is defined at the level of derived categories, it is exact, i.e., induces a functor $\text{FL}_Y : \text{Mod}_h(\mathcal{D}_{\mathcal{E}}) \rightarrow \text{Mod}_h(\mathcal{D}_{\mathcal{E}'})$.

Definition 3.8. Keep the notations of the previous definition, and assume moreover that U and U' are one-dimensional, with respective coordinates t and τ . Put $z = \tau^{-1}$, and denote by $j_\tau : \mathbb{G}_{m,\tau} \hookrightarrow \mathbb{A}_\tau^1$ and $j_z : \mathbb{G}_{m,\tau} \hookrightarrow \mathbb{A}_z^1 = \mathbb{P}_\tau^1 \setminus \{\tau = 0\}$ the canonical embeddings. Then the partial localized Fourier-Laplace transformation with respect to τ is defined by

$$\text{FL}_Y^{\text{loc}} := (j_z \times \text{id}_Y)_+ (j_\tau \times \text{id}_Y)^+ \text{FL}_Y.$$

In the sequel, we need to consider a partial compactification of the torus S_2 and an extension of φ to a projective morphism. Consider the embedding

$$(4) \quad \begin{aligned} g : S_1 &\hookrightarrow \mathbb{P}(V^\vee) = \mathbb{P}^n \\ y &\longmapsto (w_0 : \dots : w_n) = (1 : \underline{y}^{a_1} : \dots : \underline{y}^{a_n}) \end{aligned}$$

and put $X := \overline{\text{Im}(g)}$. Consider the graph $\Gamma_\varphi \subset S_1 \times \mathbb{A}^1 \times S_2$, and let ΓX be the closure of Γ_φ in $X \times \mathbb{A}^1 \times S_2$. Then we have $\Gamma X \subset Z^*$, where

$$Z^* := \mathcal{V} \left(\sum_{i=0}^n \lambda_i \cdot w_i \right) \subset \mathbb{P}^n \times \mathbb{A}^1 \times S_2$$

is the universal hypersurface. We thus have the following commutative diagram

$$(5) \quad \begin{array}{ccccc} & & k & & \\ & & \curvearrowright & & \\ & S_1 \times S_2 & \longleftrightarrow & \Gamma X & \hookrightarrow & Z^* \\ & \searrow \varphi & & \downarrow \Gamma\varphi & \swarrow p_2 & \\ & & & \mathbb{A}^1 \times S_2 & & \end{array}$$

where the map $S_1 \times S_2 \hookrightarrow \Gamma X$ is the composition of the isomorphism $S_1 \times S_2 \xrightarrow{\cong} \Gamma\varphi$ with $\Gamma\varphi \hookrightarrow \Gamma X$ and the map $\Gamma X \hookrightarrow Z^*$ is the closed embedding mentioned above.

We need also the following important geometric property of the families given by φ and ϕ .

Lemma 3.9. *The morphism $\Gamma\varphi : \Gamma X \rightarrow \mathbb{A}^1 \times S_2$ is stratified smooth on its boundary $\Gamma X \setminus (S_1 \times S_2)$. As a consequence, the restricted morphism $f_\lambda : S_1 \rightarrow \mathbb{A}^1$ is cohomologically tame in the sense of [Sab06, §8] for any parameter value $\lambda \in S_2$.*

Proof. Recall that the projective toric variety X is stratified by $X = \bigcup_{\delta \subset \Delta} X_\delta$, where Δ is the convex hull in \mathbb{R}^d of the columns of the matrix A , δ is a face of Δ , and X_δ is a torus orbit associated with the face δ (cf., for instance, [GKZ94, Prop. 5.1.9]). We obtain an induced stratification on $(\Gamma X)_\delta$ of ΓX . By definition, $\Gamma\varphi$ is stratified smooth if for each δ , the restriction $(\Gamma\varphi)|_{\Gamma X_\delta}$ to the stratum ΓX_δ is a non-singular map. One can show that $\Gamma\varphi$ is stratified smooth outside the big stratum $S_1 \times S_2$ iff the map f_λ is non-degenerate for any parameter value $\lambda \in S_2$, which holds by [RS15b, Lem. 2.8].

Fix then a $\lambda \in S_2$. The function $f_\lambda : S_1 \rightarrow \mathbb{A}^1$ is partially compactified to a projective morphism $F_\lambda : (\Gamma X)_{\mathbb{A}^1 \times \{\lambda\}} \rightarrow \mathbb{A}^1$ which is the restriction of $\Gamma\varphi$ over $\mathbb{A}^1 \times \{\lambda\}$. We have an open embedding $g_\lambda : S_1 \hookrightarrow (\Gamma X)_{\mathbb{A}^1 \times \{\lambda\}}$ (the compactification of the graph of f_λ), and f_λ is cohomologically tame if the sheaf $\mathbf{R}(g_\lambda)_* \mathbb{Q}_{S_1}$ has no vanishing cycles outside of S_1 with respect to $F_\lambda - c$ for $c \in \mathbb{C}$. As we have just proved, $\Gamma\varphi$ is stratified smooth on $\Gamma X \setminus (S_1 \times S_2)$, so that $F_\lambda - c$ is stratified smooth on $(\Gamma X)_{\mathbb{A}^1 \times \{\lambda\}} \setminus S_1$ for any $c \in \mathbb{C}$ and then the vanishing cycles with respect to it of any stratified local system, as $\mathbf{R}(g_\lambda)_* \mathbb{Q}_{S_1}$, have no support outside of S_1 by [Dim04, Prop. 4.2.8]. In conclusion, f_λ is cohomologically tame. \square

As an intermediate goal we want to compare the left-hand side of the isomorphism of Proposition 3.6 with an object derived from a certain intersection cohomology \mathcal{D} -module on Z^* .

Recall that for a smooth algebraic variety Y , and an open subvariety $j : U \hookrightarrow Y$, we call intersection cohomology module with coefficients in some \mathcal{D}_U -module \mathcal{N} is the intermediate extension $j_{\dagger+} \mathcal{N} := \text{im}(j_{\dagger} \mathcal{N} \rightarrow j_+ \mathcal{N})$. Its name comes from the fact that, if \mathcal{N} is smooth and corresponds to a local system \mathcal{L} on U under the Riemann-Hilbert correspondence, $j_{\dagger+} \mathcal{N}$ corresponds to the intersection cohomology complex on Y with coefficients in \mathcal{L} , $IC(Y, \mathcal{L})$.

We will also make use of different Radon transformations. Let us define them and state some properties, following [Rei14]:

Definition 3.10. Denote by $Z \subset \mathbb{P}(V^\vee) \times V$ the universal hyperplane given with equation $\sum_{i=0}^n w_i \lambda_i = 0$ and by $U := (\mathbb{P}(V^\vee) \times V) \setminus Z$ its complement. Consider the following diagram

$$\begin{array}{ccccc} & & U & & \\ & \swarrow \pi_1^U & \downarrow j_U & \searrow \pi_2^U & \\ \mathbb{P}(V^\vee) & \xleftarrow{\pi_1} & \mathbb{P}(V^\vee) \times V & \xrightarrow{\pi_2} & V, \\ & \swarrow \pi_1^Z & \downarrow i_Z & \searrow \pi_2^Z & \\ & & Z & & \end{array}$$

The Radon transformations are functors from $D^b(\mathcal{D}_{\mathbb{P}(V^\vee)})$ to $D^b(\mathcal{D}_V)$ given by

$$\begin{aligned}\mathcal{R} &:= \pi_{2,+}^Z \pi_1^{Z,+} \cong \pi_{2,+} i_{Z,+} i_Z^+ \pi_1^+, \\ \mathcal{R}^\circ &:= \pi_{2,+}^U \pi_1^{U,+} \cong \pi_{2,+} j_{U,+} j_U^+ \pi_1^+, \\ \mathcal{R}_c^\circ &:= \pi_{2,\dagger}^U \pi_1^{U,+} \cong \pi_{2,+} j_{U,\dagger} j_U^+ \pi_1^+, \\ \mathcal{R}_{cst} &:= \pi_{2,+} \pi_1^+.\end{aligned}$$

Proposition 3.11. *Let g be as in 4. Then, for every $\beta \in \mathbb{C}^d$, we have the following two exact sequences of regular holonomic \mathcal{D}_V -modules, which are dual to each other:*

$$\begin{aligned}0 \longrightarrow \mathcal{H}^{-1} \mathcal{R}_{cst} g_+ \mathcal{O}_{S_1}^\beta &\longrightarrow \mathcal{H}^0 \mathcal{R} g_+ \mathcal{O}_{S_1}^\beta \longrightarrow \mathcal{H}^0 \mathcal{R}_c^\circ g_+ \mathcal{O}_{S_1}^\beta \longrightarrow \mathcal{H}^0 \mathcal{R}_{cst} g_+ \mathcal{O}_{S_1}^\beta \longrightarrow 0, \\ 0 \longrightarrow \mathcal{H}^0 \mathcal{R}_{cst} g_\dagger \mathcal{O}_{S_1}^{-\beta} &\longrightarrow \mathcal{H}^0 \mathcal{R}^\circ g_\dagger \mathcal{O}_{S_1}^{-\beta} \longrightarrow \mathcal{H}^0 \mathcal{R} g_\dagger \mathcal{O}_{S_1}^{-\beta} \longrightarrow \mathcal{H}^1 \mathcal{R}_{cst} g_\dagger \mathcal{O}_{S_1}^{-\beta} \longrightarrow 0.\end{aligned}$$

Moreover, the \mathcal{D}_V -modules $\mathcal{H}^i \mathcal{R}_{cst} g_\star \mathcal{O}_{S_1}^\beta$, for $i \in \{-1, 0, 1\}$ and $\star \in \{+, \dagger\}$ that appear in the above sequences are \mathcal{O}_V -free. Consequently, for any $\beta \in \mathbb{C}^d$, calling j_{V^*} the canonical inclusion $V^* := \mathbb{A}_{\lambda_0}^1 \times S_2 \hookrightarrow V$, we have isomorphisms of $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S_2}$ -modules

$$\begin{aligned}\mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R}_c^\circ g_+ \mathcal{O}_{S_1}^\beta &\cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_+ \mathcal{O}_{S_1}^\beta \\ \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R}^\circ g_\dagger \mathcal{O}_{S_1}^\beta &\cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_\dagger \mathcal{O}_{S_1}^\beta.\end{aligned}$$

Proof. Following the notation of the previous definition, since Z is smooth, the excision triangle ([HTT08, Prop. 1.7.1]) corresponding to the diagram $U \hookrightarrow \mathbb{P}(V^\vee) \times V \leftarrow Z$ gives rise to the following triangles of Radon transformations for any $\mathcal{M} \in D_c^b(\mathcal{D}_{\mathbb{P}(V^\vee)})$:

$$\begin{aligned}(6) \quad &\mathcal{R}_{cst} \mathcal{M} \longrightarrow \mathcal{R}^\circ \mathcal{M} \longrightarrow \mathcal{R} \mathcal{M} \longrightarrow, \\ (7) \quad &\mathcal{R} \mathcal{M} \longrightarrow \mathcal{R}_c^\circ \mathcal{M} \longrightarrow \mathcal{R}_{cst} \mathcal{M} \longrightarrow,\end{aligned}$$

where the second triangle is dual to the first.

Note that $D\mathcal{O}_{S_1}^\beta \cong \mathcal{O}_{S_1}^{-1-\beta} \cong \mathcal{O}_{S_1}^{-\beta}$. It suffices then to show the existence of the first exact sequence of the statement of the proposition and the fact that the $\mathcal{H}^i \mathcal{R}_{cst} g_\star \mathcal{O}_{S_1}^\beta$ are constant. The idea of the proof is similar to that of proposition 3.6; the existence of the exact sequences here follows by just a variation of [Rei14, Prop. 2.8], but considering the twist by $y^{-1-\beta}$ of \mathcal{O}_{S_1} . That needs in turn [op. cit., Prop. 2.5, 2.7, Lem. 2.6], but almost every argument is functorial and anyway they can be easily adapted to our context. The constancy of $\mathcal{H}^i \mathcal{R}_{cst} g_\star \mathcal{O}_{S_1}^\beta$ can be proved as the second point of [op. cit., Lem. 2.7]. \square

Now is when we can properly state and prove the next comparison result of this section. Recall that we take a matrix $A \in M(d \times n, \mathbb{Z})$ satisfying the assumptions 3.1 and a parameter vector $\beta \in \mathbb{C}^d$. $Z^* = \{\sum_{i=0}^n w_i \lambda_i = 0\} \subset \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2$ denotes the universal hypersurface and the map k is

$$\begin{aligned}k : S_1 \times S_2 &\longrightarrow Z^* \\ (\underline{y}, \underline{\lambda}) &\longmapsto (1 : \underline{y}^{a_1}, \dots, \underline{y}^{a_n}, -\sum_{i=1}^n \lambda_i \underline{y}^{a_1}, \lambda_1, \dots, \lambda_n)\end{aligned}$$

Proposition 3.12. *In the above situation, we have the following isomorphism of $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S_2}$ -modules:*

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 p_{2,+} k_\dagger^+ \mathcal{O}_{S_1 \times S_2}^\beta \xrightarrow{\sim} \mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^\beta.$$

Moreover, this isomorphism is induced from the canonical morphism $k_\dagger^+ \mathcal{O}_{S_1 \times S_2}^\beta \hookrightarrow k_+ \mathcal{O}_{S_1 \times S_2}^\beta$ by applying the functor $\mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 p_{2,+}$.

Proof. Call p_1 the restriction to Z^* of the projection $\mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2 \rightarrow \mathbb{P}^n$, abusing a bit of the notation. Consider the cartesian diagram

$$\begin{array}{ccc} S_1 \times S_2 & \xrightarrow{\pi_1} & S_1 \\ k \downarrow & \square & \downarrow g \\ Z^* & \xrightarrow{p_1} & \mathbb{P}^n \end{array}$$

By base change, we have that $k_+ \mathcal{O}_{S_1 \times S_2}^\beta \cong p_1^+ g_+ \mathcal{O}_{S_1}^\beta$ for any $\beta \in \mathbb{C}^d$. Now, since p_1 is smooth, we have the analogous isomorphism $k_{\dagger} \mathcal{O}_{S_1 \times S_2}^\beta \cong p_1^+ g_{\dagger} \mathcal{O}_{S_1}^\beta$ for every β as well. Now note that because of p_1 being non-characteristic, it is easy to show (cf. the proof of the third point of [RS17, Prop. 2.22]) that for every β

$$k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^\beta \cong p_1^+ g_{\dagger+} \mathcal{O}_{S_1}^\beta.$$

Consider now the commutative diagram

$$\begin{array}{ccc} & Z^* & \xrightarrow{p_2} V^* \\ p_1 \swarrow & \downarrow j_{Z^*} & \downarrow j_{V^*} \\ \mathbb{P}^n & \xleftarrow{\pi_1^Z} Z & \xrightarrow{\pi_2^Z} V \end{array},$$

where the square is also cartesian. By applying base change again, we obtain the natural transformations of functors from $D_c^b(\mathcal{D}_{\mathbb{P}^n})$ to $D_c^b(\mathcal{D}_{V^*})$

$$(8) \quad p_{2,+} p_1^+ \cong p_{2,+} j_{Z^*}^+ \pi_1^{Z,+} \cong j_{V^*}^+ \pi_{2,+}^Z \pi_1^{Z,+} = j_{V^*}^+ \mathcal{R}.$$

Therefore, it follows that

$$(9) \quad p_{2,+} k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^\beta \cong j_{V^*}^+ \mathcal{R} g_{\dagger+} \mathcal{O}_{S_1}^\beta,$$

for any $\beta \in \mathbb{C}^d$.

We need now to relate the various Radon transformations with the Fourier-Laplace transformation $FL : \text{Mod}_h(\mathcal{D}_{V^\vee}) \rightarrow \text{Mod}_h(\mathcal{D}_V)$. This is possible due to the fundamental result of d'Agnolo and Eastwood [DE03, Prop. 1]. We quote the formulation from the proof of [RS17, Lem. 2.12]. Let $Bl_0(V^\vee) \subset \mathbb{P}^n \times V^\vee$ be the blow-up of V^\vee at the origin and consider the following diagram

$$\begin{array}{ccccc} V^* & \xrightarrow{h} & V^\vee & \xleftarrow{p} & Bl_0(V^\vee) \\ & \searrow \tilde{h} & \uparrow j_0 & & \nearrow q \\ \pi_{V^*} \downarrow & & V^\vee \setminus \{0\} & & \\ & & \downarrow \pi & & \\ S_1 & \xrightarrow{g} & \mathbb{P}^n & & \end{array}$$

where π is the canonical morphism $V^\vee \setminus \{0\} \rightarrow \mathbb{P}(V^\vee)$, π_{V^*} is the second projection, and h and \tilde{h} are given by $(y_0, \underline{y}) \mapsto (y_0, y_0 \underline{y}^{\alpha_1}, \dots, y_0 \underline{y}^{\alpha_d})$. Then we have the natural transformations

$$(10) \quad \mathcal{R}_c^\circ g_+ \cong FL h_+ \pi_{V^*}^+ \quad \text{and} \quad \mathcal{R}^\circ g_{\dagger} \cong FL h_{\dagger} \pi_{V^*}^+,$$

of functors from $\text{Mod}_h(\mathcal{D}_{S_1})$ to $\text{Mod}_h(\mathcal{D}_V)$. Indeed, although in general \mathcal{R}_c° and \mathcal{R}° are not exact, the compositions $\mathcal{R}_c^\circ g_+$ and $\mathcal{R}^\circ g_{\dagger}$ are so precisely due to the above isomorphisms.

Applying $FL_{S_2}^{\text{loc}} j_{V^*}^+$ to the natural transformations in 10 we obtain the following ones of functors from $\text{Mod}_h(\mathcal{D}_{S_1})$ to $\text{Mod}_h(\mathcal{D}_{\mathbb{A}_z^1 \times S_2})$:

$$(11) \quad FL_{S_2}^{\text{loc}} j_{V^*}^+ \mathcal{R}_c^\circ g_+ \cong FL_{S_2}^{\text{loc}} j_{V^*}^+ FL h_+ \pi^+ \quad \text{and} \quad FL_{S_2}^{\text{loc}} j_{V^*}^+ \mathcal{R}^\circ g_{\dagger} \cong FL_{S_2}^{\text{loc}} j_{V^*}^+ FL h_{\dagger} \pi^+$$

We are now closer to the isomorphism of the statement; let us rewrite in a different way the functors above to obtain it. Namely,

$$FL_{S_2}^{\text{loc}} j_{V^*}^+ FL \cong j_{V^*}^+ FL_W^{\text{loc}} FL \cong j_{V^*}^+ (j_z \times \text{id}_W)_+ (j_\tau \times \text{id}_W)^+ FL_W FL \cong j_{V^*}^+ (j_z \times \text{id}_W)_+ (j_\tau \times \text{id}_W)^+ FL_{\mathbb{A}_{w_0}^1}.$$

From the third of the assumptions 3.1 we know that we we can decompose the morphism h as

$$h : V^* \xleftarrow{h_1} \mathbb{G}_{m, w_0} \times W^\vee \xleftarrow{j_\tau \times \text{id}_{W^\vee}} \mathbb{A}_{w_0}^1 \times W^\vee = V^\vee,$$

where h_1 is a closed embedding (cf. [RS15b, Prop. 2.1 (1), proof of Thm. 2.4, p. 213]). It follows that $(j_\tau \times \text{id}_W)^+ FL_{\mathbb{A}_{w_0}^1} h_+ \cong FL_{\mathbb{A}_{w_0}^1} (j_\tau \times \text{id}_{W^\vee})^+ h_+ \cong FL_{\mathbb{A}_{w_0}^1} (j_\tau \times \text{id}_{W^\vee})^+ (j_\tau \times \text{id}_{W^\vee})_+ h_{1,+} \cong FL_{\mathbb{A}_{w_0}^1} h_{1,+}$.

Summarizing, we obtain that

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathrm{FL} h_{\star} \cong j_{V^*}^+(j_z \times \mathrm{id}_W) \mathrm{FL}_{\mathbb{A}_{w_0}^1} h_{1,+},$$

where $\star \in \{+, \dagger\}$, because $h_{1,+} = h_{1,\dagger}$ for h_1 is proper. In particular, we have $\mathrm{FL}_{S_2}^{\mathrm{loc}} \circ j_{V^*}^+ \circ \mathrm{FL} \circ h_+ \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} \circ j_{V^*}^+ \circ \mathrm{FL} \circ h_{\dagger}$. As a consequence, we can claim using the isomorphisms in 11 that

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R}_c^{\circ} g_+ \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R}^{\circ} g_{\dagger},$$

and so, from applying this last natural transformation to $\mathcal{O}_{S_1}^{\beta}$,

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_{\dagger+} \mathcal{O}_{S_1}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_+ \mathcal{O}_{S_1}^{\beta}.$$

The final step towards the statement uses the first point of [Rei14, Prop. 2.7]. Although its statement has to do only with the structure sheaf \mathcal{O}_{S_1} , as we said in the proof of Proposition 3.11 almost every argument follows by functoriality and it is straightforward to generalise them to work with $\mathcal{O}_{S_1}^{\beta}$ to get that

$$(12) \quad j_{V^*}^+ \mathcal{R} g_+ \mathcal{O}_{S_1}^{\beta} \cong \varphi_+ \mathcal{O}_{S_1 \times S_2}^{\beta}.$$

In conclusion, we finally obtain, by using the isomorphism from 9, that

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 p_{2,+} k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_{\dagger+} \mathcal{O}_{S_1}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^{\beta}.$$

The last statement is an easy consequence of isomorphisms 8, 9 and 12, noting that

$$\mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 p_{2,+} k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 j_{V^*}^+ \mathcal{R} g_{\dagger+} \mathcal{O}_{S_1}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R} g_+ \mathcal{O}_{S_1}^{\beta} \cong \mathrm{FL}_{S_2}^{\mathrm{loc}} \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^{\beta}$$

is induced from the canonical morphism $g_{\dagger+} \mathcal{O}_{S_1}^{\beta} \hookrightarrow g_+ \mathcal{O}_{S_1}^{\beta}$ via $\mathrm{FL}_{S_2}^{\mathrm{loc}} j_{V^*}^+ \mathcal{R}$. \square

Consider the space $P := \mathbb{P}^n \times (\mathbb{A}_{\lambda_0}^1 \times S_2)$, together with the canonical closed embedding $i_P : Z^* \hookrightarrow P$. We have the diagram

$$\begin{array}{ccc} Z^* & \xleftarrow{i_P} & P \\ & \downarrow p_2 & \swarrow p_2 \\ \mathbb{A}_{\lambda_0}^1 \times S_2 & & \end{array},$$

where we denote the restriction of the second projection $p_2 : P \rightarrow \mathbb{A}_{\lambda_0}^1 \times S_2$ to Z^* by the same letter (as was done before). Since i_P is proper, we have

$$\mathcal{H}^0 p_{2,+} k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \cong \mathcal{H}^0 p_{2,+} (i_P \circ k)_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \quad \text{and}$$

$$\mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^{\beta} \cong \mathcal{H}^0 p_{2,+} k_+ \mathcal{O}_{S_1 \times S_2}^{\beta} \cong \mathcal{H}^0 p_{2,+} (i_P \circ k)_+ \mathcal{O}_{S_1 \times S_2}^{\beta}.$$

so that we can also write the morphism $\mathcal{H}^0 p_{2,+} k_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \xrightarrow{\sim} \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^{\beta}$ from Proposition 3.12 as

$$(13) \quad \mathcal{H}^0 p_{2,+} (i_P \circ k)_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \longrightarrow \mathcal{H}^0 p_{2,+} (i_P \circ k)_+ \mathcal{O}_{S_1 \times S_2}^{\beta}.$$

In order to proceed, we will have to take into account certain group actions on the spaces $S_1 \times S_2$, $\mathbb{P}^n \times (\mathbb{A}_{\lambda_0}^1 \times S_2)$ and $\mathbb{A}_{\lambda_0}^1 \times S_2$ as well as some equivariance properties of the various sheaves of modules on these spaces. For the readers convenience, we recall some facts from [RS17, §2.4].

We consider the action

$$S_1 \times (S_1 \times S_2) \longrightarrow S_1 \times S_2$$

$$\underline{t}, (\underline{t}, \lambda_1, \dots, \lambda_n) \longmapsto (t_1 y_1, \dots, t_d y_d, \underline{t}^{-a_1} \lambda_1, \dots, \underline{t}^{-a_n} \lambda_n),$$

the action

$$S_1 \times (\mathbb{P}^n \times (\mathbb{A}_{\lambda_0}^1 \times S_2)) \longrightarrow (\mathbb{P}^n \times (\mathbb{A}_{\lambda_0}^1 \times S_2))$$

$$\underline{t}, (w_0 : \dots : w_n, \lambda_0, \lambda_1, \dots, \lambda_n) \longmapsto (w_0 : \underline{t}^{a_1} w_1 : \dots : \underline{t}^{a_n} w_n, \lambda_0, \underline{t}^{-a_1} \lambda_1, \dots, \underline{t}^{-a_n} \lambda_n),$$

and the action

$$\begin{aligned} S_1 \times (\mathbb{A}_{\lambda_0}^1 \times S_2) &\longrightarrow \mathbb{A}_{\lambda_0}^1 \times S_2 \\ \underline{t}, (\lambda_0, \lambda_1, \dots, \lambda_n) &\longmapsto (\lambda_0 : \underline{t}^{-a_1} \lambda_1, \dots, \underline{t}^{-a_n} \lambda_n). \end{aligned}$$

It can be shown as in loc. cit. that all these four actions are free (basically because the action $S_1 \times S_2 \rightarrow S_2$ which sends $(\underline{t}, \underline{\lambda})$ to $\underline{t}^{-a_1} \lambda_1, \dots, \underline{t}^{-a_n} \lambda_n$ is free) and hence have smooth geometric quotients. These are described by the following result, which we cite from [RS17, §2.4]. For a free action of an algebraic group G on a smooth variety X , we write X/G for the geometric quotient, which is again a smooth algebraic variety.

Proposition 3.13. *In the above situation, put $S^{\text{red}} := (\mathbb{G}_{m,t})^{n-d}$. Then the geometric quotients $(S_1 \times S_2)/S_1$, $(\mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2)/S_1$ and $(\mathbb{A}_{\lambda_0}^1 \times S_2)/S_1$ are given, respectively, by the spaces*

$$S_1 \times S^{\text{red}}, P := \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}, \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}$$

There is a canonical embedding $S^{\text{red}} \hookrightarrow S_2$ inducing embeddings (all denoted by ι)

$$\begin{aligned} S_1 \times S^{\text{red}} &\hookrightarrow S_1 \times S_2 \\ P &\hookrightarrow \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2 \\ S_1 \times \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}} &\hookrightarrow S_1 \times \mathbb{A}_{\lambda_0}^1 \times S_2. \end{aligned}$$

In the sequel, we will always consider S^{red} as a subspace of S_2 (as well as $P = \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}$ as a subspace of $\mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2$ etc.). We define reduced versions of the maps φ and $i_P \circ k$ by the cartesian diagrams

$$\begin{array}{ccc} S_1 \times S_2 & \xrightarrow{\varphi} & \mathbb{A}_{\lambda_0}^1 \times S_2 \\ \uparrow \iota & & \uparrow \iota \\ S_1 \times S^{\text{red}} & \xrightarrow{\iota' \varphi} & \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}} \end{array} \quad \text{and} \quad \begin{array}{ccc} S_1 \times S_2 & \xrightarrow{i_P \circ k} & \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2 \\ \uparrow \iota & & \uparrow \iota \\ S_1 \times S^{\text{red}} & \xrightarrow{l} & P \end{array}$$

In order to illustrate these statements, consider the main case of interest, where we have

$$A = \begin{pmatrix} 1 & -1 & 0 & 0 & \dots & 0 \\ 1 & 0 & -1 & 0 & \dots & 0 \\ \vdots & \dots & & & & \\ 1 & 0 & 0 & 0 & \dots & -1 \end{pmatrix} \in M((n-1) \times n, \mathbb{Z}),$$

Then $d = n - 1$, $S^{\text{red}} = \mathbb{G}_{m,t}$ and we have

$$(14) \quad \begin{aligned} \iota' \varphi : S_1 \times \mathbb{G}_{m,t} &\longrightarrow \mathbb{A}_{\lambda_0}^1 \times \mathbb{G}_{m,t} \\ (y_1, \dots, y_{n-1}, t) &\longmapsto \left(\frac{1}{y_1} + \dots + \frac{1}{y_{n-1}} + t \cdot y_1 \cdot \dots \cdot y_{n-1}, t \right) \end{aligned}$$

Going back to the situation of a general matrix $A \in M(d \times n, \mathbb{Z})$ satisfying the assumptions 3.1, we quote the following result from [RS17] showing that ι behaves well with respect to all modules in question. To simplify our notation we will write

$$M_{\dagger+, S_2} := \mathcal{H}^0 p_{2,+} (i_P \circ k)_{\dagger+} \mathcal{O}_{S_1 \times S_2}^\beta \quad \text{and} \quad M_{S_2} := \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times S_2}^\beta = \mathcal{H}^0 p_{2,+} (i_P \circ k)_{\dagger+} \mathcal{O}_{S_1 \times S_2}^\beta,$$

and analogously,

$$M_{\dagger+} := \mathcal{H}^0 \iota' p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^\beta \quad \text{and} \quad M := \mathcal{H}^0 \iota' \varphi_+ \mathcal{O}_{S_1 \times S^{\text{red}}}^\beta = \mathcal{H}^0 p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^\beta$$

where we write $\iota' p_2 : P = \mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}} \rightarrow \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}$ (to distinguish this projection from the one to $\mathbb{A}_{\lambda_0}^1 \times S_2$ considered above).

Proposition 3.14 ([RS17, Prop. 2.22., Lem. 6.4.]). *The morphism $\iota : \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}} \hookrightarrow \mathbb{A}_{\lambda_0}^1 \times S_2$ is non-characteristic for the modules $M_{\dagger+, S_2}$ and M_{S_2} . Consequently, from the morphism*

$$\mathcal{H}^0 p_{2,+}(i_P \circ k)_{\dagger+} \mathcal{O}_{S_1 \times S_2}^{\beta} \longrightarrow \mathcal{H}^0 \varphi_{+} \mathcal{O}_{S_1 \times S_2}^{\beta} = \mathcal{H}^0 p_{2,+}(i_P \circ k)_{+} \mathcal{O}_{S_1 \times S_2}^{\beta}$$

(see (13)) we obtain by applying ι^+ the morphism of $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}$ -modules

$$\phi : M_{\dagger+} \longrightarrow M$$

and an isomorphism of $\mathcal{D}_{\mathbb{A}_{\frac{1}{2}} \times S^{\text{red}}}$ -modules $\text{FL}_{S^{\text{red}}}^{\text{loc}} M_{\dagger+} \cong \text{FL}_{S^{\text{red}}}^{\text{loc}} M$.

Remark 3.15. From now on we will assume that the β_i are real numbers, since we will use some Hodge theoretic constructions, which are not valid for arbitrary complex β_i , as commented in Remark 3.4. We start by introducing certain filtrations on the \mathcal{D} -modules considered above.

Recall that an easy calculation decomposing $\iota^{\vee} \varphi$ as a graph embedding followed by a projection shows that the direct image complex $\iota^{\vee} \varphi_{+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ can be represented by

$$\left(\iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d}[\partial_{\lambda_0}], d - \kappa(\beta) \wedge - (df \wedge) \otimes \partial_{\lambda_0} \right),$$

where f is still the first component of $\iota^{\vee} \varphi$.

We consider the filtration on each $\iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d}[\partial_{\lambda_0}]$ given by

$$(15) \quad F_{k+l} \iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d}[\partial_{\lambda_0}] = \sum_{i=0}^{k+l} \iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d} \partial_{\lambda_0}^i.$$

One easily checks that this filtration is compatible with the differential, so that we obtain the filtered complex

$$\begin{aligned} & F_k \left(\iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d}[\partial_{\lambda_0}], d - (\kappa(\beta) \wedge) - (df \wedge) \otimes \partial_{\lambda_0} \right) := \\ & \left(F_{k+\bullet} \iota^{\vee} \varphi_{*} \Omega_{S_1 \times S^{\text{red}}/S^{\text{red}}}^{\bullet+d}[\partial_{\lambda_0}], d - (\kappa(\beta) \wedge) - (df \wedge) \otimes \partial_{\lambda_0} \right). \end{aligned}$$

Hence we get an induced filtration on the cohomology sheaves $\mathcal{H}^i \iota^{\vee} \varphi_{+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$, in particular, we have a filtration F_{\bullet} on $M = \mathcal{H}^0 \iota^{\vee} \varphi_{+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$.

On the other hand, it is well-known that $l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ carries a filtration $F^H l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ such that $(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}, F^H)$ underlies a pure polarizable complex Hodge module of weight $n = \dim(S_1) + \dim(S^{\text{red}})$ (which is by definition a filtered \mathcal{D}_P -module which is a direct summand of a pure polarizable real Hodge module (cf., e.g., [DS13, Def. 3.2.1])). Since $\iota^{\vee} p_2$ is projective (not just proper), the direct image $M_{\dagger+} = \mathcal{H}^0 \iota^{\vee} p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ also carries a Hodge filtration $F^H \mathcal{H}^0 \iota^{\vee} p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ such that $(M_{\dagger+}, F^H)$ underlies a pure polarisable complex Hodge module (see the first point of [Sai88, Thm. 1]). We consider the shifted filtration $F_{\bullet}^{H,sh} M_{\dagger+} := F_{\bullet-d}^H M_{\dagger+}$.

Lemma 3.16. *Consider again the morphism $\phi : M_{\dagger+} \rightarrow M$ from Proposition 3.14 which after applying the functor $\text{FL}_{S^{\text{red}}}^{\text{loc}}$ induces an isomorphism of $\mathcal{D}_{\mathbb{A}_{\frac{1}{2}} \times S^{\text{red}}}$ -modules. Then ϕ is filtered with respect to the filtrations on both sides, i.e., for every $k \in \mathbb{Z}$, we have the inclusion*

$$\phi \left(F_k^{H,sh} M_{\dagger+} \right) \subset F_k M.$$

Proof. Denote by $P^* = \mathbb{A}^n \times (\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}})$ is the the complement in P of the divisor $w_0 = 0$. Then the map l can be decomposed into a closed embedding $l_1 : S_1 \times S^{\text{red}} \hookrightarrow P^*$ and an open embedding $l_2 : P^* \hookrightarrow P$. Here again the fact that l_1 is closed follows from the third assumption in 3.1. We can further decompose l_1 as $l_1 = l_0 \circ i_{(\iota^{\vee} \varphi)}$, where $i_{(\iota^{\vee} \varphi)}$ is the graph embedding of $\iota^{\vee} \varphi$. More precisely, we

have the following diagram

(16)

$$\begin{array}{ccccc}
& & & & l \\
& & & & \curvearrowright \\
& & & & l_1 \\
& & & & \curvearrowright \\
S_1 \times S^{\text{red}} & \xrightarrow{i(\prime\varphi)} & S_1 \times (\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}) & \xrightarrow{l_0} & P^* = \mathbb{A}^n \times (\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}) & \xrightarrow{l_2} & P = \mathbb{P}^n \times (\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}) \\
& \searrow \prime\varphi & \searrow \prime p_2 & \searrow \prime p_2 & \searrow \prime p_2 & \searrow \prime p_2 & \searrow \prime p_2 \\
& & & & \mathbb{A}_{\lambda_0}^1 \times S^{\text{red}} & &
\end{array}$$

where we denote by slight abuse of notation all projections to the last coordinates by $\prime p_2$. Recall that $M_{\dagger+} = \mathcal{H}^0 \prime p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ and $M = \mathcal{H}^0 \prime \varphi_+ \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta} = \mathcal{H}^0 \prime p_{2,+} l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$.

Notice that the filtration $F_{\bullet}^{H_{sh}} M_{\dagger+}$ is induced from the filtration $F_{\bullet}^{H_{sh}} \text{DR}_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{\bullet+n}(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta})$ on the relative de Rham complex, where

$$(17) \quad F_k^{H_{sh}} \text{DR}_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{\bullet+n}(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) := \left(\dots \longrightarrow \Omega_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+l} \otimes F_{k+l}^{H_{sh}} l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta} \longrightarrow \dots \right)$$

by the first point of [Sai88, Thm. 1].

On the other hand, we have the isomorphism of $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}$ -modules

$$\prime p_{2,*} \text{DR}_{P^*/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \cong M$$

as the above diagram shows, just by the definition of $\mathcal{H}^0 \prime p_{2,+}$. The (shifted) Hodge filtration $F_{\bullet}^{H_{sh}}$ of the mixed Hodge module $l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ can be explicitly written down, due to the fact that l_1 is a closed embedding, and then one checks that the filtration $F_{\bullet} M$ (as defined by formula (15)) is induced from the filtration

$$(18) \quad F_k \text{DR}_{P^*/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{\bullet+n}(l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) := \left(\dots \longrightarrow \Omega_{P^*/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+l} \otimes F_{k+l}^{H_{sh}} l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta} \longrightarrow \dots \right).$$

Note that by passing to global sections on $\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}$ (using that this space is affine) the morphism $\phi : M_{\dagger+} \rightarrow M$ is induced from the following morphism of complexes

$$\begin{aligned}
R\Gamma \left(P, \text{DR}_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \right) & \xrightarrow{\phi} \Gamma \left(P^*, \text{DR}_{P^*/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \right) \\
& \cong R\Gamma \left(P^*, \text{DR}_{P^*/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{1,+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \right) \\
& \cong R\Gamma \left(P^*, \text{DR}_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \right) \\
& \cong R\Gamma \left(P^*, \text{DR}_{P/\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}^{n+\bullet}(l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}) \right)
\end{aligned}$$

Since this morphism is simply given by restricting sections of $l_{\dagger+} \mathcal{O}_{S_1 \times S^{\text{red}}}^{\beta}$ from P to P^* , we see that it is filtered with respect to the filtrations from equations (17) and (18) and so is $\phi : M_{\dagger+} \rightarrow M$ with respect to the filtrations $F_{\bullet}^{H_{sh}}$ on $M_{\dagger+}$ resp. F_{\bullet} on M . \square

In general, these two filtrations on $M_{\dagger+}$ and M are not equal, simply because the underlying $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}$ -modules are not equal. They become equal after localized partial Fourier transformation as we have shown in Proposition 3.12. First we will explain that these transformations can be performed at the filtered level and how the last result can be interpreted in this context.

We will use a general procedure which produces from a filtered $\mathcal{D}_{\mathbb{A}_{\lambda_0}^1 \times S^{\text{red}}}$ -module (N, F_\bullet) a lattice $G_0^{F_\bullet}$ inside $\text{FL}_{S^{\text{red}}}^{\text{loc}} N$, i.e., an $\mathcal{O}_{\mathbb{A}_z^1 \times S^{\text{red}}}$ -module which generates $\text{FL}_{S^{\text{red}}}^{\text{loc}} N$ over $\mathcal{D}_{\mathbb{A}_z^1 \times S^{\text{red}}}$.

Definition 3.17. (cf. [Sab08, §1.d], [SY15, §A.1]) Let X be a smooth affine variety and let (N, F_\bullet) be a filtered $\mathcal{D}_{\mathbb{A}_s^1 \times X}$ -module, which we identify with its module of global sections. Consider the algebraic microlocalization

$$N[\partial_s^{-1}] := \mathbb{C}[s]\langle \partial_s, \partial_s^{-1} \rangle \otimes_{\mathbb{C}[s]\langle \partial_s \rangle} N.$$

By letting τ act as $-\partial_s$ and ∂_τ as s , we can envision $N[\partial_s^{-1}]$ as a $\mathcal{D}_X[\tau]\langle \partial_\tau, \partial_\tau^{-1} \rangle$ -module (which actually coincides with $\text{FL}_X^{\text{loc}} N$). Let now $\widehat{\text{loc}}$ be the natural localization morphism $\widehat{\text{loc}} : N \rightarrow N[\partial_s^{-1}]$. Then we define

$$(19) \quad G_0^{F_\bullet} \text{FL}_X^{\text{loc}} N := \sum_{j \geq 0} \partial_s^{-j} \widehat{\text{loc}}(F_j N),$$

notice that then $G_0^{F_\bullet} \text{FL}_X^{\text{loc}} N$ has naturally the structure of a $\mathcal{R}_{\mathbb{A}_s^1 \times X}^{\text{int}}$ -module. We also put for any $k \in \mathbb{Z}$

$$G_k^{F_\bullet} \text{FL}_X^{\text{loc}} N := z^k \cdot G_0^{F_\bullet} \text{FL}_X^{\text{loc}} N = \sum_{j \geq 0} \partial_s^{-(j+k)} \widehat{\text{loc}}(F_j N) = \sum_{j \geq 0} \partial_s^{-j} \widehat{\text{loc}}(F_{j+k} N).$$

There is an interpretation of this construction as a Fourier-Laplace transformation for $\mathcal{R}_{\mathbb{A}_s^1 \times X}^{\text{int}}$ -modules as explained in [SY15, Rem. A.3]. Using this interpretation, one can show the following fact.

Lemma 3.18. *Let (N, F_\bullet) be a filtered $\mathcal{D}_{\mathbb{A}_s^1 \times X}$ -module underlying an element in $\text{MHM}(\mathbb{A}_s^1 \times X, \mathbb{C})$ (the abelian category of complex mixed Hodge modules). Then the $\mathcal{R}_{\mathbb{A}_z^1 \times X}^{\text{int}}$ -module $G_0^{F_\bullet} \text{FL}_X^{\text{loc}}(N)$ underlies an element of $\text{IrrMHM}(X)$.*

Proof. First we identify N with a $\mathcal{D}_{\mathbb{P}_s^1 \times X}(*\{\infty\} \times X)$ -module, so that (N, F_\bullet) underlies a Hodge module on $\mathbb{P}_s^1 \times X$. Its Rees module $\mathcal{R}_F M$ than underlies an element in $\text{MTM}^{\text{int}}(\mathbb{P}_s^1 \times X)$. Now the construction of $G_0^{F_\bullet} \text{FL}_X^{\text{loc}}(N)$ can be rephrased as the composition of the exponential twist functor $(\mathcal{T}^{s/z} \otimes -)$ (see [SY15, § 3] and [Sab15, 1.6] for more details on the exponential twist in the twistor setup) applied to $\mathcal{R}_F M$ followed by the direct image $\mathcal{H}^0 q_+$ by the second projection $q : \mathbb{P}_s^1 \times X \rightarrow X$. From [Sab15, Thm. 0.2(2)], we know that the exponential twist sends $\text{MHM}(\mathbb{P}_s^1 \times X, \mathbb{C})$ (or rather its essential image in $\text{MTM}^{\text{int}}(\mathbb{P}_s^1 \times X)$) to $\text{IrrMHM}(\mathbb{P}_s^1 \times X)$, and the projection $\mathcal{H}^0 q_+$ preserves IrrMHM according to [op. cit., Thm. 0.2(1)] since q is projective. \square

With these definitions at hand, we have the following consequence of Lemma 3.16.

Corollary 3.19. *In the above situation, we have*

$$G_0^{F_\bullet} \text{FL}_{S^{\text{red}}}^{\text{loc}} M_{\dagger+} \subset G_0^{F_\bullet} \text{FL}_{S^{\text{red}}}^{\text{loc}} M$$

Proof. This is a direct consequence of the definition in formula (19), taking into account Lemma 3.16 and the fact that the filtered morphism ϕ induces an isomorphism of $\mathcal{D}_{\mathbb{A}_z^1 \times S^{\text{red}}}$ -modules by applying the functor $\text{FL}_{S^{\text{red}}}^{\text{loc}}$. \square

From now on, we will specify the above situation to our main example, where the matrix A is given as

$$A = \begin{pmatrix} 1 & -1 & 0 & 0 & \dots & 0 \\ 1 & 0 & -1 & 0 & \dots & 0 \\ \vdots & \dots & & & & \\ 1 & 0 & 0 & 0 & \dots & -1 \end{pmatrix}$$

In particular, we have $d = n - 1$, and $S^{\text{red}} = \mathbb{G}_{m,t}$. We still write $M_{\dagger+} = \mathcal{H}^0 p_{2,+} l_{\dagger+} \mathcal{O}_{S_1 \times \mathbb{G}_{m,t}}^\beta$ and $M = \mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times \mathbb{G}_{m,t}}^\beta$. We seek to improve the inclusion of the last corollary to an equality of lattices

inside the $\mathcal{D}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}$ -modules $\mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M_{\dagger+}) \cong \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M)$. We will follow an argument from the proof of [Sab08, Lem. 4.7]. In order to do so, we have to make more explicit the structure of the module $\mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M) \cong \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M_{\dagger+})$, which is done by the next lemma.

Lemma 3.20. (1) *The singular locus $\Sigma := \mathrm{Sing}(M) = \mathrm{Sing}(\mathcal{H}^0 \varphi_+ \mathcal{O}_{S_1 \times \mathbb{G}_{m,t}}^\beta)$ is given by*

$$\Sigma := \bigcup_{\xi \in \mu_n} \{(n \cdot \xi \cdot t', t)\} \subset \mathbb{A}_{\lambda_0}^1 \times \mathbb{G}_{m,t}$$

where $(t')^n = t$ (here we chose once and for all an n -th root of t , the choice does not matter as a direct sum over all values $n\xi t'$ is taken).

- (2) *Write $D := \{0\} \times \mathbb{G}_{m,t} \subset \mathbb{A}_z^1 \times \mathbb{G}_{m,t}$. Then $\mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M)$ is $\mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}(*D)$ -locally free of rank n .*
(3) *Consider the sheaf $\widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}$ which is the formal completion of $\mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}$ along the divisor $\{0\} \times \mathbb{G}_{m,t}$. Then we have a decomposition (as sheaves on $\mathbb{G}_{m,t}$)*

$$\mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M) \otimes_{\mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}}(*D) \widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}(*D) \cong \bigoplus_{\xi \in \mu_n} \widehat{\mathcal{E}}_\xi \otimes \widehat{\mathcal{N}}_{\alpha_\xi}$$

where $\widehat{\mathcal{E}}_\xi := (\widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}(*D), d - d(n \cdot \xi \cdot t'/z))$ and $\widehat{\mathcal{N}}_{\alpha_\xi} = (\widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}(*D), d + \alpha_\xi dz/z)$, with $\alpha_\xi \in \mathbb{C}$.

- Proof.* (1) It is well known that the singular locus of a Gauss-Manin system, i.e., of the top-cohomology of the direct image complex $\varphi_+ \mathcal{M}$ is nothing but the discriminant of the morphism φ provided that the module \mathcal{M} is smooth (which is the case here, since $\mathcal{M} = \mathcal{O}_{S_1 \times \mathbb{G}_{m,t}}^\beta$). Recall (see formula (14)) that $\varphi(y_1, \dots, y_{n-1}, t) = \left(\frac{1}{y_1} + \dots + \frac{1}{y_{n-1}} + t \cdot y_1 \dots y_{n-1}, t\right)$. One easily checks (see, e.g., [DS04, §1.B.]) that a point $(y_1, \dots, y_{n-1}) \in S_1$ is critical if and only if $y_1 = \dots = y_{n-1} =: y$ and $y^n \cdot t = 1$ (and that all critical points are Morse). Then the critical values are as indicated.
(2) This is a direct consequence of the second point of [DS03, Thm. 1.11], since by the discussion above the singular locus Σ satisfies the assumption (NC) of loc. cit. It is also known that the rank of $\mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}}(M)$ equals the global Milnor number of φ , i.e., the numbers of critical points, which is n .
(3) This follows from [Sab02, Ch. III, Thm. 5.7], since the critical values of φ , i.e., the eigenvalues of the pole part of $z^2 \nabla_z$, are distinct for any $t \in \mathbb{G}_{m,t}$. □

With these preparations, we can state the next result.

Theorem 3.21. *In the above situation, we have*

$$G_0^{F \bullet H sh} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M_{\dagger+} = G_0^{F \bullet} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M.$$

Proof. We have already proved the inclusion

$$G_0^{F \bullet H sh} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M_{\dagger+} \subset G_0^{F \bullet} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M$$

of $\mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}$ -modules. Since both sheaves coincide outside the divisor $D = \{0\} \times \mathbb{G}_{m,t}$, and since $\widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}$ is $\mathcal{O}_{(\mathbb{A}_z^1 \times \mathbb{G}_{m,t}, D)}$ -flat, it is therefore sufficient to show that

$$G_0^{F \bullet H sh} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M_{\dagger+} \otimes \widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}} = G_0^{F \bullet} \mathrm{FL}_{\mathbb{G}_{m,t}}^{\mathrm{loc}} M \otimes \widehat{\mathcal{O}}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}.$$

This follows as in the proof of [Sab08, Lem. 4.7]: Using the formal decomposition result from the last Lemma, both modules can be interpreted as filtered direct images under p_2 of two modules which coincide on P^* . In these microlocal direct images, the contributions from $P \setminus P^*$ vanish by Lemma 3.9 (notice that inside $\mathbb{P}^n \times \mathbb{A}_{\lambda_0}^1 \times S_2$, we have $P^* \cap \Gamma X \cong S_1 \times S^{\mathrm{red}}$ where we see S^{red} as a subspace of S_2 via the embedding ι), and therefore both modules are equal. □

Our final result in this section is the following.

Theorem 3.22. *Let $\alpha_1, \dots, \alpha_n$ be real numbers. Let P and H be the operators*

$$P = z^2 \partial_z + ntz \partial_t + \gamma z \quad \text{and} \quad H = \prod_{i=1}^n z(t \partial_t - \alpha_i) - t,$$

where $\gamma = n\alpha_1 - \sum_{i=2}^n \alpha_i$. Consider the $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -module

$$\widehat{\mathcal{H}} := \mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}} \langle z^2 \partial_z, zt \partial_t \rangle / (P, H).$$

Then, $\widehat{\mathcal{H}}$ underlies an object of $\text{IrrMHM}(\mathbb{G}_{m,t})$, whose associated $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module is $\mathcal{H}(\alpha_i; \emptyset)$. Moreover, $\widehat{\mathcal{H}}$ can be extended in a unique way to an irreducible $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{P}^1}^{\text{int}}$ -module, $\widehat{\mathcal{H}}_{pr}$, such that it underlies an object of $\text{IrrMHM}(\mathbb{P}^1)$.

Proof. Let us assume first that $\alpha_1 = 0$, denote by α the vector $(\alpha_2, \dots, \alpha_n)$ and put $\alpha_0 := 0$. By Corollary 2.8 we know that $\mathcal{H}(\alpha_1, \dots, \alpha_n; \emptyset)$ is the inverse image $\iota^+ \mathcal{M}_A^\alpha$, where we recall that ι is the morphism $\mathbb{G}_{m,t} \rightarrow \mathbb{A}^n$ given by $t \mapsto (t, 1, \dots, 1)$ and A is the $(n-1) \times n$ -matrix of the beginning of this section. As a matter of fact, we have the same isomorphism with the twistor versions of both objects. Indeed, consider now $\iota : \mathbb{G}_{m,t} \rightarrow \mathbb{G}_m^n = S_2$. The inverse image functor in the category of twistor \mathcal{D} -modules is induced by the usual inverse image functor of \mathcal{O} -modules, $(\text{id}_{\mathbb{A}_z^1} \times \iota)^*$ in this case (cf. [Moc15a, §2.1.6.2]). Then it is easy to see that

$$\widehat{\mathcal{H}} \cong \iota^+ \widehat{\mathcal{M}}_A^{(\alpha_0, \alpha)}.$$

Namely, we replace each $z\lambda_i \partial_{\lambda_i}$ by $z\lambda_1 \partial_{\lambda_1} - z\alpha_i$, so that we can present $\widehat{\mathcal{M}}_A^{(\alpha_0, \alpha)}$ as the \mathcal{O}_{S_2} -module $\mathcal{O}_{S_2} \langle z^2 \partial_z, z\lambda_1 \partial_{\lambda_1} \rangle / \mathcal{J}$, where \mathcal{J} is generated by

$$\prod_{i=1}^n z(\lambda_1 \partial_{\lambda_1} - \alpha_i) - \lambda_1 \cdot \dots \cdot \lambda_n \quad \text{and} \quad z^2 \partial_z + nz\lambda_1 \partial_{\lambda_1} + \gamma z.$$

Now the inverse image amounts simply to set $\lambda_1 = t$ and $\lambda_i = 1$ for $i = 2, \dots, n$ in the generators of the ideal, from which the desired isomorphism follows.

From Proposition 3.6 we conclude that

$$\widehat{\mathcal{H}} \cong \iota^+ \mathcal{H}^0 \left(\pi_{2,*} \Omega_{S_1 \times S_2 / S_2}^{\bullet+d} [z], z(d - \kappa(\alpha) \wedge) - df \wedge \right) \cong \mathcal{H}^0 \left(\pi_{2,*} \Omega_{S_1 \times \mathbb{G}_{m,t} / \mathbb{G}_{m,t}}^{\bullet+d} [z], z(d - \kappa(\alpha) \wedge) - df \wedge \right),$$

and it is easy to see from Definition 3.17 that we have

$$\mathcal{H}^0 \left(\pi_{2,*} \Omega_{S_1 \times \mathbb{G}_{m,t} / \mathbb{G}_{m,t}}^{\bullet+d} [z], z(d - \kappa(\alpha) \wedge) - df \wedge \right) \cong G_0^F \bullet \text{FL}_{\mathbb{G}_{m,t}}^{\text{loc}} \mathcal{H}^0 \iota_+ \mathcal{O}_{S_1 \times \mathbb{G}_{m,t}}^\beta = G_0^F \bullet \text{FL}_{\mathbb{G}_{m,t}}^{\text{loc}} M.$$

Since by Theorem 3.21 we can further conclude

$$\widehat{\mathcal{H}} \cong G_0^F \bullet^{Hsh} \text{FL}_{\mathbb{G}_{m,t}}^{\text{loc}} M_{\uparrow+},$$

we obtain that $\widehat{\mathcal{H}}$ underlies an element of $\text{IrrMHM}(\mathbb{G}_{m,t})$ by Lemma 3.18 (recall that $M_{\uparrow+}$ underlies a pure polarizable complex Hodge module). Restricting $\widehat{\mathcal{H}}$ to $z = 1$ we get the original $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module $\mathcal{H}(\alpha_i; \emptyset)$.

Assume now that $\alpha_1 \neq 0$. The tensor product of $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -modules $\widehat{\mathcal{H}} \otimes_{\mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}} \widehat{\mathcal{K}}_{-\alpha_1}$ gives rise to the corresponding tensor product of twistor $\mathcal{D}_{\mathbb{G}_{m,t}}$ -modules. The first factor is, by the discussion above when $\alpha_1 = 0$ and [Sab15, Cor. 0.5], an irregular mixed Hodge module of exponential-Hodge origin. By loc. cit. again, since $\widehat{\mathcal{K}}_{\alpha_1}$ is the faithful image of a mixed Hodge module on $\mathbb{G}_{m,t}$, the tensor product with it preserves the condition of being in $\text{IrrMHM}(\mathbb{G}_{m,t})$, and so is the case of our original $\widehat{\mathcal{H}}$.

We will call $\widehat{\mathcal{H}}$ a classical hypergeometric $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -module, underlying a classical hypergeometric integrable twistor $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module.

Let now $j : \mathbb{G}_{m,t} \hookrightarrow \mathbb{P}^1$ be the canonical inclusion and consider the $\mathcal{D}_{\mathbb{P}^1}$ -module $\mathcal{H}_{pr} := j_{\uparrow+} \widehat{\mathcal{H}}$. It is an irreducible holonomic $\mathcal{D}_{\mathbb{P}^1}$ -module, because so is $\widehat{\mathcal{H}}$ by Proposition 2.3. Then it gives rise to

a unique pure integrable twistor $\mathcal{D}_{\mathbb{P}^1}$ -module $\widehat{\mathcal{H}}_{pr}$ by [Moc11, Thm. 1.4.4] and [Sab15, Rem. 1.40]. In addition, its underlying $\mathcal{D}_{\mathbb{P}^1}$ -module \mathcal{H}_{pr} is rigid by virtue of Proposition 2.4. As a consequence, we can invoke [Sab15, Thm. 0.7] and claim that such twistor $\mathcal{D}_{\mathbb{P}^1}$ -module is in fact an object of $\text{IrrMHM}(\mathbb{P}^1)$. Take now $\widehat{\mathcal{H}}' := j^+ \widehat{\mathcal{H}}_{pr}$, which is an irregular mixed Hodge module whose underlying $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module is \mathcal{H} , by [Moc15a, Prop. 14.1.24]. Since the functor Ξ_{DR} is faithful by [op. cit., Rem. 7.2.9], we have an injection of Hom groups

$$\text{Hom}_{\text{IrrMHM}(\mathbb{G}_{m,t})}(\widehat{\mathcal{H}}, \widehat{\mathcal{H}}') \hookrightarrow \text{Hom}_{\mathcal{D}_{\mathbb{G}_{m,t}}}(\mathcal{H}, \mathcal{H}),$$

but since \mathcal{H} is irreducible its only endomorphism is the identity, so $j^+ \widehat{\mathcal{H}}_{pr} = \widehat{\mathcal{H}}$ and we are done. \square

We end this section with a conjecture that would be the analogous result to the last theorem for arbitrary hypergeometric modules $\mathcal{H}(\alpha_i, \beta_j)$. When trying to prove it along the lines of this paper, one is faced with the problem that the matrix A that is needed to obtain $\mathcal{H}(\alpha_i, \beta_j)$ as an inverse image of a GKZ-system \mathcal{M}_A^β (see Corollary 2.8) does not satisfy the last of the assumptions 3.1, in contrast to the case $\mathcal{H}(\alpha_i, \emptyset)$. However, we believe that the conjecture will follow from the main results of [RS15a]. We plan to discuss these questions in a subsequent paper.

Conjecture 3.23. *Let $n, m \in \mathbb{N}$ and let $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_m be real numbers. Let P and H be the operators*

$$P = z^2 \partial_z + (n - m)tz \partial_t + \gamma z \quad \text{and} \quad H = \prod_{i=1}^n z(t \partial_t - \alpha_i) - t \prod_{j=1}^m z(t \partial_t - \beta_j),$$

where $\gamma = (n - m)\alpha_1 - \sum_{i=2}^n \alpha_i + \sum_{j=1}^m \beta_j$. Consider the $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -module

$$\widehat{\mathcal{H}} := \mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}} \langle z^2 \partial_z, zt \partial_t \rangle / (P, H).$$

Then, $\widehat{\mathcal{H}}$ underlies an object of $\text{IrrMHM}(\mathbb{G}_{m,t})$ with associated $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module $\mathcal{H}(\alpha_i; \beta_j)$. It can be uniquely extended to an irreducible $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{P}^1}^{\text{int}}$ -module underlying an object of $\text{IrrMHM}(\mathbb{P}^1)$.

Notice that this conjecture also covers the case of regular hypergeometric systems (i.e., the case $n = m$). Then the presence of the operator $z^2 \partial_z + \gamma z$ in the ideal defining $\widehat{\mathcal{H}}$ means precisely that $\widehat{\mathcal{H}}$ underlies an object in $\text{IrrMHM}(\mathbb{G}_{m,t})$ which belongs to the essential image of $\text{MHM}(\mathbb{G}_{m,t})$ (see the statement in [Sab15, Theorem 0.2(1)]). This is consistent with the fact that regular hypergeometric systems underly mixed Hodge modules. As stated above in Proposition 2.5, the corresponding Hodge numbers have been calculated in [Fed15], and so the conjecture is a first step towards a unifying formula for these numbers for all hypergeometric systems, including our formula (theorem 4.7 below) and Fedorov's formula.

4. THE IRREGULAR HODGE FILTRATION

We will use the following conventions and notations (cf. [Sab15, Not. 2.1]): From now on, X will mean the torus $\mathbb{G}_{m,t}$. We will write $\mathcal{X} := \mathbb{A}_z^1 \times X$, ${}^\theta \mathcal{X} = \mathcal{X} \times \mathbb{G}_{m,\theta}$, ${}^\tau \mathcal{X} = \mathcal{X} \times \mathbb{A}_\tau^1$ and ${}^\tau X_0 = X \times \{\tau = 0\}$, where $\theta = 1/\tau$. We will fix again n to be a positive integer number, and $\alpha_1, \dots, \alpha_n$ to be n real numbers, now increasingly ordered and assumed to belong to the interval $[0, 1)$. We keep on dealing with the classical hypergeometric \mathcal{D}_X -module $\mathcal{H} = \mathcal{H}(\alpha_i; \emptyset)$ and its associated twistor \mathcal{D}_X -module $\widehat{\mathcal{H}}$. We will denote the underlying $\mathcal{R}_{\mathcal{X}}^{\text{int}}$ -module by the same symbol. From [op. cit., Thm. 0.7] and from Theorem 3.22 we know that there exist a unique irregular Hodge filtration of \mathcal{H} . In this section we are going to calculate it.

For any $\mathcal{R}_{\mathcal{X}}^{\text{int}}$ -module $\widehat{\mathcal{M}}$ we can define the associated $\mathcal{R}_{{}^\theta \mathcal{X}}^{\text{int}}$ -module ${}^\theta \widehat{\mathcal{M}}$ as the inverse image $\mathcal{O}_{{}^\theta \mathcal{X}}$ -module $\mu^* \widehat{\mathcal{M}}$, endowed with a natural action of $\mathcal{R}_{{}^\theta \mathcal{X}}^{\text{int}}$ as depicted in [Sab15, 2.4] (note that $\theta = \tau^{-1}$), where μ is the morphism given in [op. cit., Not. 2.1]

$$\mu : \begin{array}{ccc} {}^\theta \mathcal{X} & \rightarrow & \mathcal{X} \\ (z, t, \theta) & \mapsto & (z\theta, t) \end{array} .$$

Proposition 4.1. *Recall that we could write $\widehat{\mathcal{H}}$ as the $\mathcal{R}_{\mathcal{X}}^{\text{int}}$ -module $\mathcal{R}_{\mathcal{X}}^{\text{int}}/(P, H)$, where P and H were, respectively,*

$$z^2\partial_z + nzt\partial_t + \gamma z \quad \text{and} \quad \prod_{i=1}^n z(t\partial_t - \alpha_i) - t,$$

for certain value of γ . Then, ${}^\theta\widehat{\mathcal{H}} = \mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(P, {}^\theta R, {}^\theta H)$, with P as before and

$${}^\theta R = z^2\partial_z - z\theta\partial_\theta \quad \text{and} \quad {}^\theta H = \prod_{i=1}^n z\theta(t\partial_t - \alpha_i) - t.$$

Proof. The morphism μ can be decomposed as $p \circ \phi$, where p is the canonical projection from ${}^\theta\mathcal{X}$ to \mathcal{X} and ϕ is the automorphism of ${}^\theta\mathcal{X}$ given by $(z, t, \theta) \mapsto (z\theta, t, \theta)$. Then, in the category of $\mathcal{O}_{\theta\mathcal{X}}$ -modules we have that

$$\mu^*\widehat{\mathcal{H}} \cong \phi^*p^*\widehat{\mathcal{H}} \cong \phi^*\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(z\theta\partial_\theta, P, H(z, t, \partial_t)) \cong \mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(z^2\partial_z - z\theta\partial_\theta, P, H(z\theta, t, \partial_t)),$$

by the chain rule.

What remains now is to prove the compatibilities of loc. cit. among the actions of $\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}$ on $\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(P, {}^\theta R, {}^\theta H)$, seen as $\mu^*\widehat{\mathcal{H}} = \mu^{-1}\widehat{\mathcal{H}} \otimes_{\mu^{-1}\mathcal{O}_{\mathcal{X}}} \mathcal{O}_{\theta\mathcal{X}}$. They are just a consequence of the presence of $z^2\partial_z - z\theta\partial_\theta$ in the ideal with which we take the quotient in $\mu^*\widehat{\mathcal{H}}$ and how z , ∂_i or ∂_z act at both sides of the tensor product. For instance, if we multiply by z at the right is the same as if we multiply by $z\theta$ at the left. \square

Remark 4.2. Let $i_{\tau=z}$ be the inclusion $\mathcal{X}^0 := \mathbb{G}_{m,z} \times X \hookrightarrow {}^\theta\mathcal{X}$ given by $(z, t) \mapsto (z, t, \tau)$. Note that, according to the fourth point of [Sab15, Lem. 2.5], we must have $i_{\tau=z}^*\widehat{\mathcal{H}} \cong \pi^{0,+}\mathcal{H}$ as $\mathcal{R}_{\mathcal{X}^0}$ -modules, with π^0 being the projection $\mathcal{X}^0 \rightarrow X$. Indeed, we have

$$\begin{aligned} i_{\tau=z}^*\widehat{\mathcal{H}} &= \mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(P, {}^\theta R, {}^\theta H, \theta z - 1) \cong \mathcal{O}_{\mathbb{G}_{m,\theta} \times \mathbb{A}_z^1 \times X} \langle zt\partial_t \rangle / (H_1, \theta z - 1) \\ &\cong \mathcal{O}_{\mathbb{G}_{m,\theta} \times \mathbb{G}_{m,z} \times X} \langle zt\partial_t \rangle / (H_1, \theta - z^{-1}) \\ &\cong \mathcal{O}_{\mathbb{G}_{m,z} \times X} \langle zt\partial_t \rangle / (H_1) \\ &\cong \mathcal{O}_{\mathbb{G}_{m,z} \times X} \otimes_{\mathcal{O}_X} \mathcal{H} \cong \pi^{0,+}\mathcal{H}, \end{aligned}$$

where H_1 is the result of replacing z by 1 at the expression of H .

In order to continue, we must pass from ${}^\theta\mathcal{X}$ to ${}^\tau\mathcal{X}$ to work in the context of [Sab15, §2.3]. Therefore, we invert θ and extend τ to the affine line to get a $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau X_0)$ -module. In other words, call $\text{inv} : \mathbb{G}_{m,\theta} \rightarrow \mathbb{G}_{m,\tau}$ the inversion operator $\theta \mapsto \theta^{-1} = \tau$ and $j : \mathbb{G}_{m,\tau} \hookrightarrow \mathbb{A}_\tau^1$ the canonical inclusion. From now on, we will denote by ${}^\tau\widehat{\mathcal{H}}$ the $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau X_0)$ -module $(\text{id}_{\mathcal{X}} \times (j \circ \text{inv}))_* \widehat{\mathcal{H}}$. By virtue of Proposition 4.1 we can write ${}^\tau\widehat{\mathcal{H}}$ as the $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau X_0)$ -module ${}^\tau\widehat{\mathcal{H}} = \mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau X_0)/(P, {}^\tau R, {}^\tau H)$, with P as before and

$${}^\tau R = z^2\partial_z + z\tau\partial_\tau \quad \text{and} \quad {}^\tau H = \prod_{i=1}^n \frac{z}{\tau} (t\partial_t - \alpha_i) - t.$$

Lemma 4.3. *For each $k = 0, \dots, n-1$, let Q_k be the operator*

$$Q_k = (-n)^k \prod_{j=1}^k \frac{z}{\tau} (t\partial_t - \alpha_j),$$

where the empty product must be understood as one. Then, the Q_k form a basis of ${}^\tau\widehat{\mathcal{H}}$ as a $\mathcal{O}_{\tau\mathcal{X}}(*^\tau X_0)$ -module. The integrable connection arising from the $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau X_0)$ -module structure associated with ${}^\tau\widehat{\mathcal{H}}$ has the following matrix expression with respect to that basis:

$$\nabla \underline{Q} = \underline{Q} \left((\tau A_0 + z A_\infty) \frac{dz}{z^2} + (-\tau A_0 + z A'_\infty) \frac{dt}{nzt} - (\tau A_0 + z A_\infty) \frac{d\tau}{z\tau} \right),$$

where A_0 , A'_∞ and A_∞ are the matrices

$$A_0 = \begin{pmatrix} 0 & & & (-n)^{nt} \\ 1 & \ddots & & 0 \\ & \ddots & 0 & \vdots \\ & & 1 & 0 \end{pmatrix}, \quad A'_\infty = \text{diag}(n\alpha_1, \dots, n\alpha_n)$$

$$\text{and } A_\infty = \text{diag}(0, 1, \dots, n-1) - \gamma I_n - A'_\infty.$$

Proof. We can again (as has been done above) use the expressions of ${}^\tau R$ and P to replace the classes of $z\tau\partial_\tau$ and $z^2\partial_z$, respectively, in terms of $zt\partial_t$. If we extend for a moment the definition of the Q_k for values of k greater than $n-1$ in the obvious way, we see that ${}^\tau\widehat{\mathcal{H}}$ is generated as a $\mathcal{O}_{\tau\mathcal{X}}(*^\tau X_0)$ -module by the Q_k , for $k \geq 0$ (since it is obviously generated by the powers of $zt\partial_t$, and those can be expressed by the Q_k 's). If we focus now at the degree in $zt\partial_t$ of the generators, we can use ${}^\tau H$ to get rid of the Q_k for $k \geq n$. Since $\deg_{zt\partial_t} Q_k = k$, they must be linearly independent over $\mathcal{O}_{\tau\mathcal{X}}(*^\tau X_0)$ and so they are a basis.

Let now $k < n-1$. Then from the relation $-nz/\tau(t\partial_t - \alpha_{k+1})Q_k = Q_{k+1}$ we can write that $nzt\partial_t Q_k = -\tau Q_{k+1} + nz\alpha_{k+1}Q_k$. Now if $k = n-1$, then

$$-nz/\tau(t\partial_t - \alpha_n)Q_{n-1} = (-n)^n \prod_{j=1}^n z/\tau(t\partial_t - \alpha_j) = (-n)^{nt}.$$

This gives us the second summand of the formula above in the statement.

The first one is a consequence of the last one and the fact that the class of $z^2\partial_z + z\tau\partial_\tau$ vanishes in ${}^\tau\widehat{\mathcal{H}}$; let us show the expression for the latter.

Take again $k < n-1$. Then,

$$z\tau\partial_\tau Q_k = (-n)^k z\tau\partial_\tau \tau^{-k} \prod_{j=1}^k z(t\partial_t - \alpha_j) = (-n)^k z(-k\tau^{-k} + \tau^{-k+1}\partial_\tau) \prod_{j=1}^k z(t\partial_t - \alpha_j) =$$

$$= -kzQ_k + Q_k z\tau\partial_\tau = -kzQ_k + Q_k z(nt\partial_t + \gamma) = z(nt\partial_t + \gamma - k)Q_k = -\tau Q_{k+1} + z(n\alpha_{k+1} + \gamma - k)Q_k.$$

The analogous calculation for $k = n-1$ gives us that $z\tau\partial_\tau Q_{n-1} = -\tau(-n)^{nt} + z(n\alpha_n + \gamma - (n-1))Q_{n-1}$. \square

Definition 4.4. For each $\alpha \in \mathbb{R}$, let us define the following subsets of ${}^\tau\widehat{\mathcal{H}}$:

$$\begin{aligned} \mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}} &:= \left\{ \sum_{k=0}^{n-1} f_k \tau^{\nu_k} Q_k : f_k \in \mathcal{O}_{\mathcal{X}}[\tau], \max(k - n\alpha_{k+1} - \gamma - \nu_k) \leq \alpha \right\}, \\ \mathcal{T}U_{<\alpha} {}^\tau\widehat{\mathcal{H}} &:= \left\{ \sum_{k=0}^{n-1} f_k \tau^{\nu_k} Q_k : f_k \in \mathcal{O}_{\mathcal{X}}[\tau], \max(k - n\alpha_{k+1} - \gamma - \nu_k) < \alpha \right\}. \end{aligned}$$

Remark 4.5. Note that the $\mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}}$ form an increasing filtration of ${}^\tau\widehat{\mathcal{H}}$, indexed by the real numbers but with a discrete set of jumping numbers, such that $\tau \mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}} = \mathcal{T}U_{\alpha-1} {}^\tau\widehat{\mathcal{H}}$ and $z\partial_\tau \mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}} \subseteq \mathcal{T}U_{\alpha+1} {}^\tau\widehat{\mathcal{H}}$ for any α . The graded piece associated with α is $\text{Gr}_\alpha^{\mathcal{T}U} {}^\tau\widehat{\mathcal{H}} = \mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}} / \mathcal{T}U_{<\alpha} {}^\tau\widehat{\mathcal{H}}$.

In the definition of the $\mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}}$ all the exponents ν_k of the powers of τ accompanying the $f_k Q_k$ satisfy that $\nu_k \geq -\alpha + k - n\alpha_{k+1} - \gamma$. Therefore, we can define the steps of the filtration in an alternative way, as the free $\mathcal{O}_{\mathcal{X}}[\tau]$ -modules of finite rank

$$\mathcal{T}U_\alpha {}^\tau\widehat{\mathcal{H}} = \bigoplus_{k=0}^{n-1} \mathcal{O}_{\mathcal{X}}[\tau] \cdot \tau^{\nu_\alpha(k)} Q_k,$$

where $\nu_\alpha(k) = \lceil -\alpha + k - \gamma - n\alpha_{k+1} \rceil$.

With this expression it is easy to see that $\tau U_\alpha \widehat{\mathcal{H}}/(\tau - z)\tau U_\alpha \widehat{\mathcal{H}}$ is the z -graded free \mathcal{O}_X -module $\bigoplus_k \mathcal{O}_X z^{\nu_\alpha(k)} \bar{Q}_k$, where

$$\bar{Q}_k = (-n)^k \prod_{j=1}^k (t\partial_t - \alpha_j).$$

As in the case of \mathcal{D} -modules, there is the notion of strict specializability and V -filtration for \mathcal{R}^{int} -modules; check [Moc15a, §2.1.2] for more details. In the setting under consideration we will use τV to denote the canonical V -filtration of a $\mathcal{R}_{\tau X}^{\text{int}}$ -module, to make clear the variety over which we are working (note the same convention in [Sab15], from Remark 2.20 on).

Proposition 4.6. *$\widehat{\mathcal{H}}$ is strictly \mathbb{R} -specializable along τX_0 and its τV -filtration is in fact given by the $\tau U_\alpha \widehat{\mathcal{H}}$.*

Proof. First of all we will see that $\tau U_\alpha \widehat{\mathcal{H}}$ is the τV -filtration of $\widehat{\mathcal{H}}$, following [Moc15a, §§2.1.2.1, 2.1.2.2]. Apart from what we already shown at the remark above, from the alternative expression of the filtration we can see that the graded pieces are $\mathcal{O}_{\mathbb{A}_z^1}$ -flat and thus strict. What remains then is showing condition (v) of [op. cit., §2.1.2.1]. Consider then the mappings $\mathfrak{p}, \mathfrak{e}$ given by

$$\begin{aligned} (\mathfrak{p}, \mathfrak{e}) : \mathbb{R} \times \mathbb{C} &\longrightarrow \mathbb{R} \times \mathbb{C} \\ (\beta, \omega) &\longmapsto (\beta + 2\Re(z\bar{\omega}), -\beta z + \omega - \bar{\omega}z^2) \end{aligned}$$

We ought to see now that the operator $z\tau\partial_\tau - \mathfrak{e}(\beta, \omega)$ is nilpotent on the graded pieces $\text{Gr}_\alpha^{\tau U} \widehat{\mathcal{H}}$ only for a finite amount of $(\beta, \omega) \in \mathcal{K} := \{\beta + 2\Re(z_0\bar{\omega}) = \alpha\}$, for any value z_0 of z . Moreover, $\widehat{\mathcal{H}}$ will be strictly \mathbb{R} -specializable if those (β, ω) belong in fact to $\mathbb{R} \times \{0\}$ (cf. [Sab15, §1.3.a]).

Take then $(\beta, \omega) \in \mathcal{K}$ and $f\tau^\nu Q_k \in \tau U_\alpha \widehat{\mathcal{H}}$, with $f \in \mathcal{O}_X[\tau]$. We must have that $k - n\alpha_{k+1} - \gamma - \nu \leq \alpha$. Assume that $k < n - 1$. Thanks to Lemma 4.3 we know that

$$(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))f\tau^\nu Q_k = (z\tau\partial_\tau + (\nu + n\alpha_{k+1} + \gamma - k + \beta)z - \omega + \bar{\omega}z^2)(f)\tau^\nu Q_k - \tau^{\nu+1}Q_{k+1}.$$

Recall that the α_i are increasingly ordered. Thus $f\tau^{\nu+1}Q_{k+1}$ lives in $\tau U_\alpha \widehat{\mathcal{H}}$, for

$$k + 1 - n\alpha_{k+2} - \gamma - \nu - 1 \leq ((k + 1) - n\alpha_{k+2} - \gamma) - (k - n\alpha_{k+1} - \gamma) - 1 + \alpha \leq \alpha.$$

Now we should look at what happens to the class of $f\tau^{\nu+1}Q_{k+1}$ in the α -graded piece of $\widehat{\mathcal{H}}$.

Note that $[f\tau^\nu Q_k] \neq 0$ if and only if $\nu + n\alpha_{k+1} + \gamma - k + \alpha = 0$, so

$$\begin{aligned} (z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))f\tau^\nu Q_k &= (z\tau\partial_\tau + (\beta - \alpha)z - \omega + \bar{\omega}z^2)(f)\tau^\nu Q_k - \tau^{\nu+1}Q_{k+1} = \\ &= (z\tau\partial_\tau - 2\Re(z_0\bar{\omega})z - \omega + \bar{\omega}z^2)(f)\tau^\nu Q_k - \tau^{\nu+1}Q_{k+1}. \end{aligned}$$

Now notice that τ divides $\tau\partial_\tau(f)$, so in fact $z\tau\partial_\tau(f)\tau^\nu Q_k \in \tau U_{\alpha-1} \widehat{\mathcal{H}}$ and then we can further reduce our expression to

$$(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))f\tau^\nu Q_k = (-\omega - 2\Re(z_0\bar{\omega})z + \bar{\omega}z^2)f\tau^\nu Q_k - \tau^{\nu+1}Q_{k+1}.$$

On the other hand, $\tau^{\nu+1}Q_{k+1}$ does not vanish either in $\text{Gr}_\alpha^{\tau U} \widehat{\mathcal{H}}$ if and only if $\alpha_{k+2} = \alpha_{k+1}$. Indeed, we know that $\nu + n\alpha_{k+1} + \gamma - k + \alpha = 0$, so doing the same as before, $k + 1 - n\alpha_{k+2} - \gamma - \nu - 1 = \alpha + n(\alpha_{k+2} - \alpha_{k+1})$ and the claim follows. Furthermore, in order to $(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))$ to vanish, we should impose that $\omega = 0$, just by looking at the coefficients of the powers of z in the expression for f .

Now if $k = n - 1$, then everything would be the same as before except $-\tau^{\nu+1}Q_{k+1}$, which becomes $-(-n)^n t\tau^{\nu+1}$, whose class vanishes obviously in the graded piece under consideration.

In conclusion, $(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))^l f\tau^\nu Q_k$ can only vanish in $\text{Gr}_\alpha^{\tau U} \widehat{\mathcal{H}}$ if $\alpha = \beta$ (and then $\omega = 0$), and does not do so until we get to an index $k + l$ such that α_{k+l} is strictly bigger than α_k . Since there is a finite set of indexes, $(z\tau\partial_\tau + \alpha z)$ is nilpotent, of nilpotency index n at most. Summing up and noting that all the calculation was in fact independent of z_0 , $\widehat{\mathcal{H}}$ is strictly \mathbb{R} -specializable along τX_0 . \square

Theorem 4.7. *Let as before $\alpha_1, \dots, \alpha_n$ be real numbers in $[0, 1)$ and put $\mathcal{H} = \mathcal{H}(\alpha_i, \emptyset)$. For each $k = 1, \dots, n$, set $\rho(k) = -n\alpha_k + k$. Then the jumping numbers of the irregular Hodge filtration of \mathcal{H} are, up to an overall real shift, the numbers $\rho(k)$. The irregular Hodge numbers are the multiplicities of those jumping numbers, or in other words, the nonzero values of $|\rho^{-1}(x)|$, for x real.*

Notice that this result has a similar shape to Proposition 2.5 which treats the case of regular hypergeometric systems. We hope that an extension of our methods will lead to a computation of irregular Hodge numbers for arbitrary irregular systems $\mathcal{H}(\alpha_i; \beta_j)$ (see again Conjecture 3.23). Such a general formula, if it exists, could then certainly be compared to the result in the regular case.

Proof. Since we know that $\widehat{\mathcal{H}}$ underlies an object in $\text{IrrMHM}(\mathbb{G}_{m,t})$ by Theorem 3.22, we conclude by [Sab15, Def. 2.52] that $\widehat{\mathcal{H}}$ is well-rescalable (as defined in [op. cit., Def. 2.19]) and so we apply [op. cit., Def. 2.22]. From Remark 4.5, we have

$$i_{\tau=z}^* \tau V_{\alpha} \tau \widehat{\mathcal{H}} = \tau V_{\alpha} \tau \widehat{\mathcal{H}} / (\tau - z) \tau V_{\alpha} \tau \widehat{\mathcal{H}} = \bigoplus_k \mathcal{O}_{\mathcal{X}} z^{\nu_{\alpha}(k)} \bar{Q}_k,$$

which is z -graded of finite rank, so the z -adic filtration on $\pi^* \mathcal{H}[z^{-1}]$ induces a filtration on $i_{\tau=z}^* \tau V_{\alpha} \tau \widehat{\mathcal{H}}$, given by

$$F_r i_{\tau=z}^* \tau V_{\alpha} \tau \widehat{\mathcal{H}} := \bigoplus_{s \leq r} \left(\bigoplus_{k: s \geq \nu_{\alpha}(k)} \mathcal{O}_{\mathcal{X}} \bar{Q}_k \right) z^s.$$

Then, $\text{Gr}^F \left(i_{\tau=z}^* \tau V_{\alpha} \tau \widehat{\mathcal{H}} \right)$ is the Rees module associated to a new good filtration $F_{\alpha+}^{\text{irr}}$ on \mathcal{H} , for some $k = 0, \dots, n-1$, which is the irregular Hodge filtration. More concretely, $F_{\bullet}^{\text{irr}} \mathcal{H}$ is given by

$$F_{\alpha+j}^{\text{irr}} = \bigoplus_{k: j \geq \nu_{\alpha}(k)} \mathcal{O}_{\mathcal{X}} \bar{Q}_k.$$

Therefore, its jumping numbers are $-\gamma + j - 1 - n\alpha_j$ for $j = 1, \dots, n$. Since the irregular Hodge filtration is defined up to an overall real shift, we can normalize the jumping numbers to $j - n\alpha_j$ and the irregular Hodge numbers will be their multiplicities. \square

REFERENCES

- [Ari10] D. Arinkin, *Rigid irregular connections on \mathbb{P}^1* , Compos. Math. **146** (2010), no. 5, 1323–1338.
- [BZMW13] Christine Berkesch Zamaere, Laura Felicia Matusevich, and Uli Walther, *Torus equivariant \mathcal{D} -modules and hypergeometric systems*, preprint arXiv:1308.5901 [math.AG], 2013.
- [DE03] Andrea D’Agnolo and Michael Eastwood, *Radon and Fourier transforms for \mathcal{D} -modules*, Adv. Math. **180** (2003), no. 2, 452–485.
- [Dim04] Alexandru Dimca, *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, 2004.
- [DS03] Antoine Douai and Claude Sabbah, *Gauss-Manin systems, Brieskorn lattices and Frobenius structures. I*, Ann. Inst. Fourier (Grenoble) **53** (2003), no. 4, 1055–1116.
- [DS04] ———, *Gauss-Manin systems, Brieskorn lattices and Frobenius structures. II*, Frobenius manifolds, Aspects Math., E36, Vieweg, Wiesbaden, 2004, pp. 1–18.
- [DS13] Michael Dettweiler and Claude Sabbah, *Hodge theory of the middle convolution*, Publ. Res. Inst. Math. Sci. **49** (2013), no. 4, 761–800.
- [ESY17] Hélène Esnault, Claude Sabbah, and Jeng-Daw Yu, *E_1 -degeneration of the irregular Hodge filtration*, J. Reine Angew. Math. **729** (2017), 171–227.
- [Fed15] Roman Fedorov, *Variations of Hodge structures for hypergeometric differential operators and parabolic Higgs bundles*, preprint arXiv:1505.01704 [math.AG], 2015.
- [GKZ94] Israel M. Gel’fand, Mikhail M. Kapranov, and Andrei V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1994.
- [GMS09] Ignacio de Gregorio, David Mond, and Christian Sevenheck, *Linear free divisors and Frobenius manifolds*, Compositio Mathematica **145** (2009), no. 5, 1305–1350.
- [HTT08] Ryoshi Hotta, Kiyoshi Takeuchi, and Toshiyuki Tanisaki, *\mathcal{D} -modules, perverse sheaves, and representation theory*, Progress in Mathematics, vol. 236, Birkhäuser Boston, Inc., Boston, MA, 2008, Translated from the 1995 Japanese edition by Takeuchi.

- [Kat90] Nicholas M. Katz, *Exponential sums and differential equations*, Annals of Mathematics Studies, vol. 124, Princeton University Press, Princeton, NJ, 1990.
- [KKP17] Ludmil Katzarkov, Maxim Kontsevich, and Tony Pantev, *Bogomolov-Tian-Todorov theorems for Landau-Ginzburg models*, J. Differential Geom. **105** (2017), no. 1, 55–117.
- [Moc11] Takuro Mochizuki, *Wild harmonic bundles and wild pure twistor \mathcal{D} -modules*, Astérisque (2011), no. 340, x+607.
- [Moc15a] ———, *Mixed twistor \mathcal{D} -modules*, Lecture Notes in Mathematics, vol. 2125, Springer, Cham, 2015.
- [Moc15b] ———, *Twistor property of GKZ-hypergeometric systems*, preprint arXiv:1501.04146 [math.AG], 2015.
- [Rei14] Thomas Reichelt, *Laurent Polynomials, GKZ-hypergeometric Systems and Mixed Hodge Modules*, Compositio Mathematica (**150**) (2014), 911–941.
- [RS15a] Thomas Reichelt and Christian Sevenheck, *Hypergeometric Hodge modules*, preprint arXiv:1503.01004 [math.AG], 2015.
- [RS15b] ———, *Logarithmic Frobenius manifolds, hypergeometric systems and quantum \mathcal{D} -modules*, Journal of Algebraic Geometry **24** (2015), no. 2, 201–281.
- [RS17] ———, *Non-affine Landau-Ginzburg models and intersection cohomology*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 3, 665–753.
- [Sab02] Claude Sabbah, *Déformations isomonodromiques et variétés de Frobenius*, Savoirs Actuels, EDP Sciences, Les Ulis, 2002, Mathématiques.
- [Sab06] ———, *Hypergeometric periods for a tame polynomial*, Port. Math. (N.S.) **63** (2006), no. 2, 173–226.
- [Sab08] ———, *Fourier-Laplace transform of a variation of polarized complex Hodge structure.*, J. Reine Angew. Math. **621** (2008), 123–158.
- [Sab15] ———, *Irregular Hodge theory*, (with the collaboration of Jeng-Daw Yu), preprint arXiv:1511.00176 [math.AG], 2015.
- [Sai88] Morihiko Saito, *Modules de Hodge polarisables*, Publ. Res. Inst. Math. Sci. **24** (1988), no. 6, 849–995 (1989).
- [Sai90] ———, *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. **26** (1990), no. 2, 221–333.
- [Sch73] Wilfried Schmid, *Variation of Hodge structure: the singularities of the period mapping*, Invent. Math. **22** (1973), 211–319.
- [Sha17] Yota Shamoto, *Hodge-Tate conditions for Landau-Ginzburg models*, preprint arXiv:1709.03244 [math.AG], 2017.
- [Sim90] Carlos T. Simpson, *Harmonic bundles on noncompact curves*, J. Amer. Math. Soc. **3** (1990), no. 3, 713–770.
- [SY15] Claude Sabbah and Jeng-Daw Yu, *On the irregular Hodge filtration of exponentially twisted mixed Hodge modules*, Forum Math. Sigma **3** (2015), e9, 71.
- [Yu14] Jeng-Daw Yu, *Irregular Hodge filtration on twisted de Rham cohomology*, Manuscripta Math. **144** (2014), no. 1-2, 99–133.

E-mail address: `alcas@hrz.tu-chemnitz.de`

E-mail address: `christian.sevenheck@mathematik.tu-chemnitz.de`

FAKULTÄT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT CHEMNITZ. 09107 CHEMNITZ (GERMANY)