

On the b -Functions of Hypergeometric Systems

Thomas Reichelt¹, Christian Sevenheck², and Uli Walther^{3,*}

¹Mathematisches Institut, Universität Heidelberg, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany, ²Fakultät für Mathematik, Technische Universität Chemnitz, 09107 Chemnitz, Germany, and ³Department of Mathematics, Purdue University, 150 N. University St., West Lafayette, IN 47907, USA

*Correspondence to be sent to: e-mail: walther@math.purdue.edu

For any integer $d \times (n + 1)$ matrix A and parameter $\beta \in \mathbb{C}^d$ let $M_A(\beta)$ be the associated A -hypergeometric (or GKZ) system in the variables x_0, \dots, x_n . We describe bounds for the (roots of the) b -functions of both $M_A(\beta)$ and its Fourier transform along the hyperplanes ($x_j = 0$). We also give an estimate for the b -function for restricting $M_A(\beta)$ to a generic point.

1 Introduction

Let D be the ring of algebraic \mathbb{C} -linear differential operators on \mathbb{C}^{n+1} with coordinates x_0, \dots, x_n .

Definition 1.1 (Compare [4, 5]). Let M be a left D -module and pick an element $m \in M$ with annihilator $I \subseteq D$. If $(V^i D)$ is the vector space spanned by the monomials $x^\alpha \partial^\beta$ with $\alpha_0 - \beta_0 \geq i$ then the b -function of $m \in M$ along the coordinate hyperplane $x_0 = 0$ is the minimal monic polynomial $b(s)$ that satisfies: $b(x_0 \partial_0) \cdot m \in (V^1 D) \cdot m$ in M , which is to say $b(x_0 \partial_0) \in I + (V^1 D)$ in D .

Received July 14, 2016; Revised February 5, 2017; Accepted February 8, 2017
Communicated by Prof. Maskai Kashiwara

If M is cyclic, that is, $M = D/I$, then we call b -function of M the b -function in the above sense of the element $1 + I \in M$. \square

The b -function exists in greater generality along any hypersurface ($f = 0$), as long as the module M is holonomic, cf. [4]. The V -filtration of Kashiwara and Malgrange then takes the form $(V^i D) = \{P \in D \mid f^{i+k} \text{ divides } P \bullet f^k \text{ for } k \gg 0\}$. Both the V -filtration and the b -function are intimately connected to the restriction of the given D -module to the hypersurface. The purpose of this note is to give, for any A -hypergeometric system as well as its Fourier transform, an explicit arithmetic description of a bound for the root set of the b -function along any coordinate hyperplane that involves the parameter β in a very elementary way.

We have several applications in mind: first, it is a longstanding question to understand the monodromy of A -hypergeometric systems, and for this purpose the roots of the b -function can be of some use. On the other hand, the Fourier transform of an A -hypergeometric system often (see [15]) appears as a direct image module under a natural torus embedding given by the columns of the matrix A . This point of view turns out to be extremely useful for Hodge theoretic considerations of A -hypergeometric systems (see [9]). It is one of the fundamental insights of Morihiko Saito (see [11, Section 3.2]) that the boundary behavior of variations of Hodge structures (or, more generally, of mixed Hodge modules) is controlled by the Kashiwara–Malgrange filtration along such a boundary divisor. In the case of a cyclic D -module, such as A -hypergeometric systems or their Fourier transforms, one can often deduce a large part of this filtration from the values of the b -function. We refer to [10] for an immediate application of our results. In a third direction, one can also see our calculation of the b -function of the Fourier transform as a refinement of [1, 15] geared towards restriction of A -hypergeometric systems.

In the last part, we compute an upper bound for the b -function of restriction of the A -hypergeometric system to a generic point, again in elementary terms of A and β . Since the restriction of a D -module to a point is a dual object to the zeroth level solution functor, our estimate can be viewed as a step towards a sheafification in β of the solution space, a problem that remains unsolved.

2 Basic notions and results

Notation. Throughout, the base field is \mathbb{C} and we consider a \mathbb{C} -vector space V of dimension $n + 1$. \square

In this introductory section, we review basic facts on A -hypergeometric systems as well as the Euler–Koszul functor. Readers are advised to refer to [6] for more detailed explanations.

Notation 2.1. For any integer matrix A , let R_A (respectively O_A) be the polynomial ring over \mathbb{C} generated by the variables ∂_j (respectively x_j) corresponding to the columns \mathbf{a}_j of A . We identify O_A with the symmetric algebra on $\text{Hom}_{\mathbb{C}}(V, \mathbb{C}) \cong \bigoplus \mathbb{C} \cdot x_j$. Further, let D_A be the ring of \mathbb{C} -linear differential operators on O_A , where we identify $\frac{\partial}{\partial x_j}$ with ∂_j and multiplication by x_j with x_j so that both R_A and O_A become subrings of D_A . \square

2.1 A -hypergeometric systems

Let $A = (\mathbf{a}_0, \dots, \mathbf{a}_n)$ be an integer $d \times (n + 1)$ matrix, $d \leq n + 1$. For convenience, we assume that $\mathbb{Z}A = \mathbb{Z}^d$. For $(v_1, \dots, v_r) = \mathbf{v} \in \mathbb{Z}^r$, we denote by $\mathbf{v}_+, \mathbf{v}_-$ the vectors given by

$$(\mathbf{v}_+)_j = \max(0, v_j) \quad \text{and} \quad (\mathbf{v}_-)_j = \max(0, -v_j).$$

For the complex parameter vector $\beta \in \mathbb{C}^d$ consider the system of d *homogeneity equations*

$$E_i \bullet \phi = \beta_i \cdot \phi, \tag{2.1}$$

where $E_i = \sum_{j=0}^n a_{i,j} x_j \partial_j$ is the i -th *Euler operator*, together with the *toric* (partial differential) *equations*

$$\{(\underbrace{\partial^{\mathbf{v}_+} - \partial^{\mathbf{v}_-}}_{:=\Delta_{\mathbf{v}}}) \bullet \phi = 0 \mid A \cdot \mathbf{v} = 0\}. \tag{2.2}$$

In R_A , the toric operators $\{\Delta_{\mathbf{v}} \mid A \cdot \mathbf{v} = 0\}$ generate the *toric ideal* I_A . The quotient

$$S_A := R_A / I_A$$

is naturally isomorphic to the semigroup ring $\mathbb{C}[\text{NA}]$. In D_A , the left ideal generated by all equations (2.1) and (2.2) is the *hypergeometric ideal* $H_A(\beta)$. We put

$$M_A(\beta) := D_A / H_A(\beta);$$

this is the A -hypergeometric system introduced and first investigated by Gelfand, Graev, Kapranov, and Zelevinsky, in [2] and a string of other articles. \diamond

2.2 A -degrees

If the rowspan of A contains $\mathbf{1}_A$ we call A *homogeneous*. Homogeneity is equivalent to I_A defining a projective variety, and also to the system $H_A(\beta)$ having only regular singularities [3, 13]. A more general A -degree function on R_A and D_A is induced by:

$$-\deg_A(x_j) := \mathbf{a}_j =: \deg_A(\partial_j).$$

We denote $\deg_{A,i}(-)$ the A -degree function associated to the weight given by the i -th row of A , so $\deg_A = (\deg_{A,1}, \dots, \deg_{A,d})$.

An R_A - (respectively D_A -)module M is A -graded if it has a decomposition $M = \bigoplus_{\alpha \in \mathbb{Z}^d} M_\alpha$ such that the module structure respects the grading $\deg_A(-)$ on R_A (respectively D_A) and M . If N is an A -graded R_A -module, then we denote $\deg_A(N) \subseteq \mathbb{Z}^d$ the set of all degrees of all non-zero homogeneous elements of N . The *quasi-degrees* $\text{qdeg}_A(N)$ of N are the points in the Zariski closure in \mathbb{C}^d of $\deg_A(N)$.

As is common, if M is A -graded then $M(\mathbf{b})$ denotes for each $\mathbf{b} \in \mathbb{Z}^d$ its shift with graded structure $(M(\mathbf{b}))_{b'} = M_{\mathbf{b}+\mathbf{b}'}$.

2.3 Euler–Koszul complex

Since

$$\begin{aligned} x^u E_i - E_i x^u &= -(A \cdot \mathbf{u})_i x^u, \\ \partial^u E_i - E_i \partial^u &= (A \cdot \mathbf{u})_i \partial^u, \end{aligned}$$

we have

$$E_i P = P(E_i - \deg_{A,i}(P)) \tag{2.3}$$

for any A -homogeneous $P \in D_A$ and all i .

On the A -graded D_A -module M one can thus define commuting D_A -linear endomorphisms E_i via

$$E_i \circ m := (E_i + \deg_{A,i}(m)) \cdot m$$

for A -homogeneous elements $m \in M$. In particular, if N is an A -graded R_A -module one obtains commuting sets of D_A -endomorphisms on the left D_A -module $D_A \otimes_{R_A} N$ by

$$E_i \circ (P \otimes Q) := (E_i + \deg_{A,i}(P) + \deg_{A,i}(Q))P \otimes Q.$$

The *Euler–Koszul complex* $\mathcal{K}_\bullet(N; \beta)$ of the A -graded R_A -module N is the homological Koszul complex induced by $E - \beta := \{(E_i - \beta_i) \circ\}_1^d$ on $D_A \otimes_{R_A} N$. In particular, the terminal module $D_A \otimes_{R_A} N$ sits in homological degree zero. We denote the homology groups of $\mathcal{K}_\bullet(N; \beta)$ by $\mathcal{H}_\bullet(N; \beta)$. Implicit in the notation is “ A ”: different presentations of semigroup rings that act on N yield different Euler–Koszul complexes.

If $N(\mathbf{b})$ denotes the usual shift-of-degree functor on the category of graded R_A -modules, then $\mathcal{K}_\bullet(N; \beta)(\mathbf{b})$ and $\mathcal{K}_\bullet(N(\mathbf{b}); \beta - \mathbf{b})$ are identical.

2.4 The toric category

There is a bijection between faces τ of the cone $\mathbb{R}_{\geq 0}A$ and A -graded prime ideals $I_A^\tau = I_A + R_A\{\partial_j \mid j \notin \tau\}$ of R_A containing I_A . If the origin is a face of $\mathbb{R}_{\geq 0}A$, it corresponds to the ideal $I_A^\emptyset = (\partial_0, \dots, \partial_n)$. In general, $R_A/I_A^\tau \cong \mathbb{C}[\mathbb{N}\tau]$.

An R_A -module N is *toric* if it is A -graded and has a (finite) A -graded composition chain

$$0 = N_0 \subsetneq N_1 \subsetneq N_2 \cdots \subsetneq N_k = N$$

such that each composition factor N_i/N_{i-1} is isomorphic as A -graded R_A -module to an A -graded shift $(R_A/I_A^\tau)(\mathbf{b})$ for some $\mathbf{b} \in \mathbb{Z}A$ and some face τ . The category of toric modules is closed under the formation of subquotients and extensions.

For toric input N , the modules $\mathcal{K}_\bullet(N; \beta)$ are holonomic. As D_A is R_A -free, any short exact sequence $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ of A -graded R_A -modules produces a long exact sequence of Euler–Koszul homology. If β is not a quasi-degree of N then the complex $\mathcal{K}_\bullet(N; \beta)$ is exact, and if N is a maximal Cohen–Macaulay module then $\mathcal{K}_\bullet(N; \beta)$ is a resolution of $\mathcal{H}_0(N; \beta)$.

2.5 The Euler space

Notation 2.2. The \mathbb{C} -linear span of the Euler operators $\{E_i\}_1^d$ is called the *Euler space*. Let E be in the Euler space. Then E is in a unique fashion (as $\text{rk}(A) = d$) a linear combination $E = \sum c_i E_i$. With $\beta_E := \sum c_i \beta_i$, we have $E - \beta_E \in H_A(\beta)$. We further write $\deg_E(-)$ for the degree function $\sum c_i \deg_{\beta_i}(-)$. \square

Denote $\theta_j = x_j \partial_j$ and $\theta = (\theta_0, \dots, \theta_n)$. A linear combination $\sum_j v_j \theta_j$ is in the Euler space if and only if the coefficient vector $\mathbf{v} = (v_0, \dots, v_n)$, interpreted as a linear functional on \mathbb{C}^{n+1} via $\mathbf{v}((q_0, \dots, q_n)) := \sum v_i q_i$, is the pull-back via A of a linear functional

on \mathbb{C}^d . In other words,

$$[\mathbf{v} \cdot \theta^T = \sum_j v_j \theta_j \text{ is in the Euler space}] \Leftrightarrow [\mathbf{v} = \mathbf{c} \cdot A \text{ for some } \mathbf{c} \in \mathbb{C}^d].$$

If $L: \mathbb{C}^d \rightarrow \mathbb{C}$ is a linear functional then the Euler operator in $H_A(\beta)$ corresponding to its image under $\text{Hom}_{\mathbb{C}}(\mathbb{C}^d, \mathbb{C}) \xrightarrow{A} \text{Hom}_{\mathbb{C}}(\mathbb{C}^{n+1}, \mathbb{C})$ is denoted $E_L - \beta_L$.

Lemma 2.3. For any set F of columns of A contained in a hyperplane that passes through the origin of \mathbb{C}^d but does not contain \mathbf{a}_k , there is an Euler operator $E_F - \beta_F$ in $H_A(\beta)$ such that the coefficient of θ_j in E_F is zero for all $j \in F$, and equal to 1 for $j = k$. If $\mathbb{R}_{\geq 0}F$ is a facet of $\mathbb{R}_{\geq 0}A$ then $E_F - \beta_F$ is unique. \square

Proof. Choose for any such set F a linear functional $L: \mathbb{Q}^d \rightarrow \mathbb{Q}$ that vanishes on F while $L(\mathbf{a}_k) = 1$. The corresponding Euler operator $E_L - \beta_L$ has the desired properties, and if we define numbers $a_{L,j}$ by

$$E_L =: \sum_j a_{L,j} x_j \partial_j$$

then $a_{L,j} = L(\mathbf{a}_j)$. The uniqueness in the facet case is obvious. \blacksquare

3 Restricting the Fourier transform

The Fourier transform $\mathcal{F}(-)$ is a functor from the category of D -modules on V to the category of D -modules on the dual space $V^* = \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$. In this section, we bound the b -function along a coordinate hyperplane of the Fourier transform $\mathcal{F}(M_A(\beta))$ of the hypergeometric system. This module is called \check{M}_A^β in [10].

The square of the Fourier transform is the involution induced by $x \mapsto -x$, which has no effect on the analytic properties of the modules we study. In particular, b -functions along coordinate hyperplanes are unaffected by this involution and we therefore consider $\mathcal{F}^{-1}(M_A(\beta))$ without harm.

We start with introducing some notation.

Notation 3.1. Let $\{y_j\}$ be the coordinates on V^* such that $\mathcal{F}^{-1}(\partial_j) = y_j$ on the level of differential operators. We let \tilde{D}_A be the ring of \mathbb{C} -linear differential operators on $\tilde{O}_A := \mathbb{C}[y_0, \dots, y_n]$, generated by $\{y_j, \delta_j\}_0^n$ where δ_j denotes $\frac{\partial}{\partial y_j}$. Then $\mathcal{F}^{-1}(x_j) = -\delta_j$. The subring $\mathbb{C}[\delta_1, \dots, \delta_n]$ of \tilde{D}_A is denoted \tilde{R}_A . The isomorphism $(\tilde{-}): D_A \rightarrow \tilde{D}_A$ induced by $\tilde{\partial}_j := y_j$ and $\tilde{x}_j = \delta_j$ sends O_A to \tilde{R}_A and R_A to \tilde{O}_A .

Thus, $\tilde{I}_A := \mathcal{F}^{-1}(I_A)$ is an ideal of \tilde{O}_A ; the advantage of considering \mathcal{F}^{-1} rather than \mathcal{F} is that \tilde{I}_A retains the shape of the generators of I_A as differences of monomials. For each j set $\tilde{\theta}_j := \mathcal{F}^{-1}(\theta_j) = -\delta_j \gamma_j$. The i -th level V -filtration on \tilde{D}_A along γ_t is spanned by $\delta^\alpha \gamma^\beta$ with $\beta_t - \alpha_t \geq i$. \square

Before, we get into the technical part, let us show by example an outline of what is to happen.

Example 3.2. Let $A = \begin{pmatrix} -1 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}$, a matrix whose associated semigroup ring is a normal complete intersection. We will estimate the b -function for restriction to the hyperplane $\gamma_1 = 0$ (corresponding to the middle column) of $\mathcal{F}^{-1}(M_A(\beta))$.

The ideal $\tilde{H}_A(\beta) := \mathcal{F}^{-1}(H_A(\beta))$ is generated by

$$-\tilde{\theta}_0 + \tilde{\theta}_2 - \beta_1, \quad \tilde{\theta}_0 + \tilde{\theta}_1 + \tilde{\theta}_2 - \beta_2, \quad \gamma_0 \gamma_2 - \gamma_1^2. \quad (3.1)$$

Since $\gamma_1 \in (V^1 \tilde{D}_A)$, $\gamma_0 \gamma_2$ and hence also $\tilde{\theta}_0 \tilde{\theta}_2$ are in $(V^1 \tilde{D}_A) + \tilde{H}_A(\beta)$. The strategy of the example, and of the theorem in this section, is to multiply the element $1 \in \tilde{D}_A$ by suitable Euler operators so that the result is a sum of a polynomial $p(\tilde{\theta}_1)$ with an element of $\mathbb{C}[\tilde{\theta}_0, \tilde{\theta}_1, \tilde{\theta}_2] \cdot \tilde{\theta}_0 \tilde{\theta}_2$; this certifies $p(\tilde{\theta}_1)$ to be in $\tilde{H}_A(\beta) + (V^1 \tilde{D}_A)$.

In the case at hand, the relevant Euler operators are $2\tilde{\theta}_0 + \tilde{\theta}_1 + \beta_1 - \beta_2$ and $\tilde{\theta}_1 + 2\tilde{\theta}_2 - \beta_1 - \beta_2$. Modulo $\tilde{H}_A(\beta)$ we can rewrite $(V^1 \tilde{D}_A) \ni 4\delta_0 \delta_2 \gamma_1^2 \equiv 4\tilde{\theta}_0 \tilde{\theta}_2 \equiv (-\tilde{\theta}_1 - \beta_1 + \beta_2)(-\tilde{\theta}_1 + \beta_1 + \beta_2)$. It follows that $(\tilde{s} + \beta_1 - \beta_2)(\tilde{s} - \beta_1 - \beta_2)$ is a multiple of the b -function, where $\tilde{s} = \tilde{\theta}_1 = -\gamma_1 \delta_1 - 1$. This Fourier twist in the argument of the b -function occurs naturally throughout and we will make our computations in this section in terms of $b(\tilde{s})$.

The expressions $\tilde{\theta}_1 + 2\tilde{\theta}_2$ and $2\tilde{\theta}_0 + \tilde{\theta}_1$ that appear in the Euler operators we used can be found systematically as follows. Let d_1, d_2 denote the coordinates on the degree group \mathbb{Z}^2 corresponding to E_1 and E_2 ; compare the discussion following Notation 2.2. An element of S_A has degree on the facet $\mathbb{R}_{\geq 0} \mathbf{a}_0$ if and only if the functional $L_1(d_1, d_2) = d_1 + d_2$ vanishes, and the Euler field that corresponds to this functional in the spirit of Lemma 2.3 is exactly $\theta_1 + 2\theta_2 - \beta_1 - \beta_2$. The elements of S_A with degree on the facet $\mathbb{R}_{\geq 0} \mathbf{a}_2$ are determined by the vanishing of $L_2(d_1, d_2) = d_2 - d_1$ and the Euler field corresponding to this functional is exactly $2\theta_0 + \theta_1 + \beta_1 - \beta_2$. It is no coincidence that the union of the kernels of these two functionals is exactly the set of quasi-degrees of $S_A / \partial_1 \cdot S_A$. The point is that modulo $\tilde{H}_A(\beta)$ all monomials in \tilde{S}_A with degree in $\mathbb{R}_+ A$ are already in $(V^1 \tilde{D}_A)$. The task is then to deal with those with degree on the boundary through multiplication with suitable expressions.




Fig. 1. Restriction of the Fourier transform to $y_1 = 0$.

Figure 1 shows as dots the elements of $\mathbb{N} A$ (where those inside A are shaded at height one); the quasi-degrees of $S_A/\partial_1 \cdot S_A$ are on the two indicated lines. Finally, $(\beta_2 - \beta_1)\mathbf{a}_1$ and $(\beta_1 + \beta_2)\mathbf{a}_1$ are the intersections of $\mathbb{R} \cdot \mathbf{a}_1$ with $\text{qdeg}_A(S_A) + \beta$. \square

We now generalize the computation of the example to the general case.

Convention 3.3. For the remainder of this section, we consider restriction to the hyperplane y_0 in order to save overhead (in terms of a further index variable). \square

Consider the toric module $N = S_A/\partial_0 S_A$, and take a toric filtration

$$(N) \quad 0 = N_0 \subsetneq N_1 \subsetneq \dots \subsetneq N_k = N$$

with composition factors

$$\overline{N}_\alpha := N_\alpha/N_{\alpha-1},$$

each isomorphic to some shifted face ring $S_{F'_\alpha}(\mathbf{b}_\alpha)$, $F'_\alpha = \tau_\alpha \cap A$, attached to a face τ_α of $\mathbb{R}_{\geq 0}A$. (We will call such F'_α also a face.) Lifting the N_α to S_A yields an increasing sequence of A -graded ideals $J_\alpha \ni \partial_0$ of S_A with $N_\alpha = J_\alpha/\partial_0 \cdot S_A$.

Choose for each composition factor a facet F_α containing F'_α . None of the faces F'_α will contain \mathbf{a}_0 (as ∂_0 is zero on N but not nilpotent on any face ring of a face containing \mathbf{a}_0) and hence we can arrange that the corresponding facets do not contain \mathbf{a}_0 either.

Lemma 2.3 produces for each \overline{N}_α a facet F_α and corresponding functional L_{F_α} (which we abbreviate to L_α) that vanishes on the facet and evaluates to 1 on \mathbf{a}_0 . The associated Euler operator in $H_A(\beta)$ is $E_{F_\alpha} - \beta_{F_\alpha}$. Since L_α is zero on all A -columns in F_α and since \overline{N}_α is a shifted quotient of S_{F_α} , there is a unique value for L_α on the A -degrees of

all non-zero A -homogeneous elements of \overline{N}_α . We denote this value by $L_\alpha(\overline{N}_\alpha)$. However, that $L_\alpha(\overline{N}_\alpha)$ does very much depend on the choice of the facet F_α even though the notation does not remember this.

Now let T_α be the image in $\mathcal{F}^{-1}(M_A(\beta))$ of $\mathcal{F}^{-1}(J_\alpha)$ under the map induced by $\tilde{O}_A \longrightarrow \tilde{D}_A \longrightarrow \mathcal{F}^{-1}(M_A(\beta))$. The image of $T_0 = y_0 \tilde{O}_A$ in $\mathcal{F}^{-1}(M_A(\beta))$ is in $(V^1 \tilde{D}_A) \cdot \bar{1}$, the bar denoting cosets in $\mathcal{F}^{-1}(M_A(\beta))$.

Lemma 3.4. In the context of the three preceding paragraphs, let κ_α be the constant $L_\alpha(\overline{N}_\alpha)$. Then in $\mathcal{F}^{-1}(M_A(\beta))$, modulo the image of $(V^1 \tilde{D}_A)$,

$$(\tilde{\theta}_0 + \kappa_\alpha - \beta_\alpha) \cdot (V^0 \tilde{D}_A) \cdot T_\alpha = (V^0 \tilde{D}_A) \cdot (\tilde{\theta}_0 + \kappa_\alpha - \beta_\alpha) \cdot T_\alpha \subseteq (V^0 \tilde{D}_A) \cdot T_{\alpha-1}. \quad \square$$

Proof. Since the commutators $[\tilde{\theta}_0, (V^0 \tilde{D}_A)]$ are in $(V^1 \tilde{D}_A)$, it suffices to show that $(\tilde{\theta}_0 + \kappa_\alpha - \beta_\alpha) \cdot T_\alpha \subseteq (V^0 \tilde{D}_A) \cdot T_{\alpha-1}$ modulo $\mathcal{F}^{-1}(H_A(\beta))$.

By definition, $\tilde{E}_\alpha - \beta_\alpha := \mathcal{F}^{-1}(E_\alpha - \beta_\alpha)$ is zero in $\mathcal{F}^{-1}(M_A(\beta))$. Take a monomial $\tilde{m} \in \tilde{O}_A$ whose coset lies in $T_\alpha \setminus T_{\alpha-1}$. By Equation (2.3), $\tilde{E}_\alpha \cdot \tilde{m} = \tilde{m}(\tilde{E}_\alpha - \kappa_\alpha)$ since $\mathcal{F}^{-1}(-)$ is a homomorphism. Now write $E_\alpha = \sum a_{\alpha,j} \theta_j$; as before, we have $a_{\alpha,j} = L_\alpha(\mathbf{a}_j)$.

Since the coefficient of θ_0 in E_α is 1, it follows that in $\mathcal{F}^{-1}(M_A(\beta))$:

$$\begin{aligned} \tilde{\theta}_0 \tilde{m} &= (-\tilde{E}_\alpha + \tilde{\theta}_0) \tilde{m} + \tilde{E}_\alpha \tilde{m} \\ &= \sum_{\substack{j \neq 0 \\ L_\alpha(\mathbf{a}_j) \neq 0}} a_{\alpha,j} \delta_j y_j \tilde{m} + \tilde{m}(\tilde{E}_\alpha - \kappa_\alpha) \\ &= \sum_{\substack{j \neq 0 \\ \mathbf{a}_j \notin F_\alpha}} a_{\alpha,j} \delta_j y_j \tilde{m} + \tilde{m}(\beta_\alpha - \kappa_\alpha). \end{aligned}$$

Recall that F_α contains F'_α and that \overline{N}_α is a $\mathbb{Z}A$ -shift of $S_{F'_\alpha} = R_A/I_{A'}^t$, whence each y_j with $\mathbf{a}_j \notin F'$ annihilates $\mathcal{F}^{-1}(\overline{N}_\alpha)$. Therefore, each term $a_{\alpha,j} \delta_j (y_j \tilde{m})$ in the last sum of the display is in $(V^0 D_A) T_{\alpha-1}$. It follows that in $\mathcal{F}^{-1}(M_A(\beta))$, we have $(\tilde{\theta}_0 + \kappa_\alpha - \beta_\alpha) T_\alpha \subseteq (V^0 \tilde{D}_A) T_{\alpha-1}$ as claimed. \blacksquare

Theorem 3.5. For $t = 0, \dots, n$, the number $\varepsilon \in \mathbb{C}$ is a root of the b -function $b(\tilde{s})$ (with $\tilde{s} = \tilde{\theta}_t = -\delta_t y_t$) of $\mathcal{F}^{-1}(M_A(\beta))$ along $y_t = 0$, only if $\varepsilon \cdot \mathbf{a}_t$ is a point of intersection of the line $\mathbb{C} \cdot \mathbf{a}_t$ with the set $\beta - \text{qdeg}_A(N)$, the quasi-degrees of the toric module $N = S_A / \partial_t S_A$ multiplied by -1 and shifted by β . \square

Proof. Without loss of generality we shall suppose that $t = 0$ by way of re-indexing.

We will show that a divisor of $\prod_{\alpha}(\tilde{\theta}_0 + \kappa_{\alpha} - \beta_{\alpha})$ is inside $H_A(\beta) + (V^1\tilde{D}_A)$, in notation from the previous lemma.

Indeed, it follows from Lemma 3.4 that $\prod_{\alpha}(\tilde{\theta}_0 + \kappa_{\alpha} - \beta_{\alpha})$ multiplies $\bar{1} \in \mathcal{F}^{-1}(M_A(\beta))$ into $(V^0\tilde{D}_A) \cdot \gamma_0 \cdot \bar{1} \subseteq (V^1\tilde{D}_A) \cdot \bar{1}$. Hence the root set of the b -function $b(\tilde{\theta}_0)$ in question is a subset of $\{\beta_{\alpha} - \kappa_{\alpha}\}$, α running through the indices of the chosen composition series of N . This set is determined by the composition series (N) and the choices of the facets F_{α} for each N_{α} . Varying over all choices of facets $\{F_{\alpha}\}$ for a given chain (N) , the root set of $b(\tilde{\theta}_0)$ is in the intersection ρ_N of all possible sets $\{\beta_{\alpha} - \kappa_{\alpha}\}_{\alpha \in (N)}$.

Since $L_{\alpha}(\mathbf{a}_0) = 1$, the point $(\beta_{\alpha} - \kappa_{\alpha}) \cdot \mathbf{a}_0$ is the intersection of the hyperplane $L_{\alpha} = \beta_{\alpha} - \kappa_{\alpha}$ with the line $\mathbb{C} \cdot \mathbf{a}_0$. Thus, ρ_N is inside the intersection of $\mathbb{C} \cdot \mathbf{a}_0$ with all arrangements $\text{Var} \prod_{\alpha}(L_{\alpha} - \beta_{\alpha} + \kappa_{\alpha})$. The intersection of the arrangements $\text{Var} \prod_{\alpha}(L_{\alpha} - \beta_{\alpha} + \kappa_{\alpha})$ is the union of the quasi-degrees of all \bar{N}_{α} of the composition chain (N) , multiplied by -1 and shifted by $-\beta_{\alpha}$. As N is finitely generated, $\text{qdeg}_A(N) = \bigcup_{\alpha} \text{qdeg}_A(\bar{N}_{\alpha})$. Hence the root set of $b(\tilde{\theta}_0)$ is contained in the intersection $-\text{qdeg}_A(S_A/\partial_0 S_A) + \beta$ with $\mathbb{C} \cdot \mathbf{a}_0$. ■

Remark 3.6. The quantity $\tilde{\theta}_t$ is the more natural argument for the b -function here. The roots of $b(\gamma_t \delta_t)$ are those of $b(\tilde{\theta}_t)$ shifted up by 1 and then multiplied by -1 . □

Example 3.7. Let $A = (\mathbf{a}_0, \mathbf{a}_1, \mathbf{a}_2) = \begin{pmatrix} -1 & 0 & 3 \\ 1 & 1 & 1 \end{pmatrix}$ and $\beta = \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}$. The ring S_A is a complete intersection but not normal.

Consider restriction to $\gamma_1 = 0$ (the middle column). Then $N = S_A/\partial_1 \cdot S_A$ has a toric filtration involving four steps, given by the ideals $0 \subsetneq \partial_0^3 \cdot N \subsetneq \partial_0^2 \cdot N \subsetneq \partial_0 \cdot N \subsetneq N$. The corresponding A -graded composition factors are $S_A(-3 \cdot \mathbf{a}_0)/(\partial_1, \partial_2)S_A$ and $\{S_A(-\alpha \cdot \mathbf{a}_0)/(\partial_0, \partial_1)S_A\}_{\alpha=0}^2$. The b -function $b(\tilde{\theta}_1)$ for the inverse Fourier transform is $(\tilde{\theta}_1 - \beta_1 - \beta_2) \prod_{\alpha=0}^2 (\tilde{\theta}_1 - \frac{3\beta_2 - \beta_1 - 4\alpha}{3})$.

Explicitly, $\gamma_1^4 - \gamma_0^3 \gamma_2 \in \tilde{H}_A(\beta)$ gives $(V^1\tilde{D}_A) \ni \delta_0^3 \delta_2 \gamma_0^3 \gamma_2 = \tilde{\theta}_2 \tilde{\theta}_0 (\tilde{\theta}_0 - 1) (\tilde{\theta}_0 - 2)$ which modulo $\tilde{H}_A(\beta)$ equals $(-1)^4 (\tilde{\theta}_1 - \beta_1 - \beta_2) \prod_{\alpha=0}^2 (\tilde{\theta}_1 - \frac{3\beta_2 - \beta_1 - 4\alpha}{3})$. The relevant Euler operators are $\theta_1 + 4\theta_2 - \beta_1 - \beta_2$ and $3\theta_1 + 4\theta_0 - 3\beta_2 + \beta_1$.

Figure 2 shows in the elements of $\mathbb{N}A$ (where those inside A are shaded at height one); the quasi-degrees of $N = S_A/\partial_1 \cdot S_A$ are on the indicated four lines. The roots of $b(\delta_1 \gamma_1)$ (which are opposite to the roots of $b(\tilde{\theta}_1)$) are the intersections of the line $\mathbb{C} \cdot \binom{0}{1}$ with the shift of the indicated lines by $-\beta$.

In this example, each composition factor corresponds to a facet and to a component of the quasi-degrees of N . One checks that each composition chain must have these




Fig. 2. Restriction of the Fourier transform to $y_1 = 0$.

four lines as quasi-degrees. However, that composition chains are far from unique and in general such correspondence will not exist. \square

Remark 3.8. The b -function for $\mathcal{F}^{-1}(M_A(\beta))$ along a coordinate hyperplane is generally not reduced, and its degree may be lower than the length of the shortest toric filtration for $N = S_A/\partial_t \cdot S_A$ would suggest. (Not every component of $\beta - \text{qdeg}_A(N)$ needs to meet the line $\mathbb{C} \cdot \mathbf{a}_t$). \square

Corollary 3.9. The roots of the b -function $b(\delta_t Y_t)$ of $\mathcal{F}^{-1}(M_A(\beta))$ along $y_t = 0$ are in the field $\mathbb{Q}(\beta)$.

Consider $\mathcal{F}^{-1}(M_A(0))$; then:

1. the roots of the b -function $b(\tilde{\theta}_t)$ are non-negative rationals;
2. if S_A is normal, all roots are in the interval $[0, 1)$;
3. if the interior ideal of S_A is contained in $\partial_t \cdot S_A$ then zero is the only root. \square

Proof. The first claim is a consequence of the intersection property in Theorem 3.5: the defining equations for the quasi-degrees are rational.

Let $N = S_A/\partial_t S_A$. For items 1.-3., we need to study the intersection of $\text{qdeg}_A(N)$ with $\mathbb{C} \cdot \mathbf{a}_t$, since $\beta = 0$ and $\delta_t Y_t = -\tilde{\theta}_t$. The quasi-degrees of N are covered by hyperplanes of the sort $L_\alpha = \varepsilon$ where L_α is a rational supporting functional of the facet F_α . In particular, we can arrange L_α to be zero on F_α , positive on the rest of A , and $L_\alpha(\mathbf{a}_t) = 1$. As $\text{deg}_A(N) \subseteq \text{deg}_A(S_A)$, $\varepsilon \geq 0$. Hence $\text{Var}(L_\alpha - \varepsilon)$ meets $\mathbb{C} \cdot \mathbf{a}_t$ in the non-negative rational multiple $\varepsilon \mathbf{a}_t$ of \mathbf{a}_t . If S_A is normal, $\text{deg}_A(S_A/\partial_t S_A)$ is covered by hyperplanes $\text{Var}(L_\alpha - \varepsilon)$ that do not meet the cone $\mathbf{a}_t + \mathbb{R}_{\geq 0} A$. These are precisely the ones for which $\varepsilon < 1$.

If $\partial_t \cdot S_A$ contains the interior ideal then $\text{deg}_A(N)$, and hence $\text{qdeg}_A(N)$, is inside the supporting hyperplanes of the cone, which meet $\mathbb{C} \cdot \mathbf{a}_t$ at the origin. \blacksquare

Remark 3.10. One special case in which case 3 of Corollary 3.9 applies is when S_A is Gorenstein and where further ∂_t generates the canonical module. The matrix $A =$

$$(\mathbf{a}_0, \dots, \mathbf{a}_3) = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

with the interior ideal being generated by $\partial_1 \partial_3$, provides an example that case (3) can occur in a Gorenstein situation without the boundary of $\mathbb{N}A$ being saturated. See [14] for a discussion on Cohen–Macaulayness of face rings of Cohen–Macaulay semigroup rings. \square

4 b -functions for the hypergeometric system

4.1 Restriction along a hyperplane

We are here interested in the b -function for the hypergeometric module $M_A(\beta)$ along the hyperplane $x_t = 0$. As in the previous section, apart from examples, we actually carry out all computations for $t = 0$, in order to have as few variables around as possible. On the other hand, the natural argument for expressing the b -function will be $s = x_0 \partial_0$.

Notation 4.1. With $A = (\mathbf{a}_0, \dots, \mathbf{a}_n)$ and distinguished index 0, we denote $A' := (\mathbf{a}_1, \dots, \mathbf{a}_n)$. Via $\mathbb{N}A' \subseteq \mathbb{N}A$ we consider $S_{A'}$ as a subring of S_A .

For $k \in \mathbb{N}$ let $\bar{J}_{A,0;k} \subseteq S_{A'}$ be the vector space spanned by the monomials $\partial^{\mathbf{u}}$ with $u_0 = 0$ (so that $\partial^{\mathbf{u}} \in S_{A'}$) that satisfy $\partial_0^k \cdot \partial^{\mathbf{u}} \in S_{A'}$. We denote $J_{A,0;k} \subseteq R_{A'}$ the preimage of $\bar{J}_{A,0;k}$ under the natural surjection $R_{A'} \rightarrow S_{A'}$. Put $J_{A,0} = \sum_{k \geq 1} J_{A,0;k}$ and $\bar{J}_{A,0} = J_{A,0}/I_{A'} \subseteq S_{A'}$. \square

Each $\bar{J}_{A,0;k}$ is a monomial ideal of $S_{A'}$ since $\partial_0^k(\partial^{\mathbf{v}} \partial^{\mathbf{u}}) = \partial^{\mathbf{v}}(\partial_0^k \partial^{\mathbf{u}})$. Note, however, that $\bar{J}_{A,0;k}$ need not be contained in $\bar{J}_{A,0;k+1}$. If $\mathbf{a}_0 \in \mathbb{R}_{\geq 0}A'$ then some power of ∂_0 is in $S_{A'}$ and so $\bar{J}_{A,0} = S_{A'}$.

Definition 4.2. For $\mathbf{a}_0 \in \mathbb{R}^d$ outside $\mathbb{R}_{\geq 0}A'$, a point $\mathbf{a} \in \mathbb{R}_{\geq 0}A'$ is \mathbf{a}_0 -visible if $\mathbf{a} + \lambda \cdot \mathbf{a}_0$, $0 < \lambda \ll 1$ is outside $\mathbb{R}_{\geq 0}A'$. (The idea behind the choice of language is that the observer stands at the point of projective space given by the line $\mathbb{R}\mathbf{a}_0$.)

By abuse of notation, we say that $\partial^{\mathbf{a}}$ is \mathbf{a}_0 -visible if \mathbf{a} is. \square

Lemma 4.3. Assume that \mathbf{a}_0 is not in the cone $\mathbb{R}_{\geq 0}A'$. Then the radical of $J_{A,0}$ is generated by the \mathbf{a}_0 -invisible elements of $S_{A'}$, and in consequence the quasi-degrees of $S_{A'}/J_{A,0}$ are a union of shifted face spans where each face is in its entirety visible from \mathbf{a}_0 . \square

Proof. If $\mathbb{Z}A/\mathbb{Z}A'$ has positive rank then all points of $\mathbb{N}A$ are \mathbf{a}_0 -visible while $J_{A,0}$ is clearly zero, so that in this case there is nothing to prove. We therefore assume that $\mathbb{Z}A/\mathbb{Z}A'$ is finite.

It is immediate that \mathbf{a} is \mathbf{a}_0 -visible if and only if any positive integer multiple of it is. This implies that no power of an \mathbf{a}_0 -visible element $\partial^{\mathbf{a}}$ of $S_{A'}$ can be in the radical of $J_{A,0}$ since $\partial^{m\mathbf{a}+k\mathbf{a}_0}$ can't have its degree in the cone of A' .

For the converse, suppose \mathbf{a} is not \mathbf{a}_0 -visible, so that there are positive integers $p < q$ with $\mathbf{a} + (p/q) \cdot \mathbf{a}_0 \in \mathbb{R}_{\geq 0}A'$. Then a high power of $\partial^{q\mathbf{a}+p\mathbf{a}_0}$ is in $\mathbb{C}[\mathbb{Z}A \cap \mathbb{R}_{\geq 0}A']$ and a suitable power $\partial^{\mathbf{b}}$ of that will be in $\mathbb{C}[\mathbb{Z}A' \cap \mathbb{R}_{\geq 0}A']$ because of the finiteness of $\mathbb{Z}A/\mathbb{Z}A'$. Now let τ be the smallest face of $\mathbb{R}_{\geq 0}A'$ that contains \mathbf{b} ; this makes \mathbf{b} an interior point of τ . Since $\mathbb{C}[\tau \cap \mathbb{Z}A']$ is a finitely generated $\mathbb{C}[\tau \cap \mathbb{N}A']$ -module, some power of $\partial^{\mathbf{b}}$ is in $\mathbb{C}[\tau \cap \mathbb{N}A'] \subseteq S_{A'}$. This shows that some power of $\partial^{q\mathbf{a}}$ times some power of $\partial^{p\mathbf{a}_0}$ is in $S_{A'}$, establishing the first claim of the lemma.

In every composition chain for $S_{A'}/J_{A,0}$, each composition factor is an $S_{A'}/\sqrt{J_{A,0}}$ -module. Thus the quasi-degrees of $S_{A'}/J_{A,0}$ are inside a union of shifted quasi-degrees of $S_{A'}/\sqrt{J_{A,0}}$ and hence all \mathbf{a}_0 -visible, which implies the second claim. \blacksquare

Our main theorem in this section is:

Theorem 4.4. The root locus of the b -function $b(x_0\partial_0)$ for restriction of $M_A(\beta)$ along $x_0 = 0$ is, up to inclusion of non-negative integers, contained in the locus of intersection $(-\text{qdeg}_{A'}(S_{A'}/\bar{J}_{A,0}) + \beta) \cap \mathbb{C} \cdot \mathbf{a}_0$. The set of integers needed can be taken to be the integers $0, \dots, k-1$ such that $J_{A,0} = \sum_{1 \leq i \leq k} J_{A,0;i}$.

In two extreme cases, one can be explicit:

1. if $\dim S_A - 1 = \dim S_{A'}$ then the b -function is linear with root given by the intersection of $(-\text{qdeg}_A(S_{A'}) + \beta) \cap \mathbb{C} \cdot \mathbf{a}_0$;
2. if $\mathbf{a}_0 \in \mathbb{R}_{\geq 0}A'$ then the b -function has integer roots in $\{0, 1, \dots, k-1\}$, where $k = \min\{t \in \mathbb{N} \mid 0 \neq t \cdot \mathbf{a}_0 \in \mathbb{N}A'\}$. \square

Proof. We first dispose of the extreme cases. If $\dim S_A - 1 = \dim S_{A'}$, then S_A is the polynomial ring $S_{A'}[\partial_0]$ and A' is a facet of A . By Lemma 2.3 there is $\mathbf{v} = (v_1, \dots, v_d)$ such that the Euler operator

$$E - \beta_E = \sum v_i(E_i - \beta_i)$$

is in $H_A(\beta)$ and equals $\theta_0 - \beta_E$. In particular, the b -function is $s - \beta_E$. On the other hand: $\bar{J}_{A,0}$ is zero in this case, $\mathbf{v} = (v_1, \dots, v_d)$ is in the kernel of A'^T , and $\mathbf{a}_0^T \mathbf{v} = 1$. Therefore, the quasi-degrees of $S_{A'}/\bar{J}_{A,0}$ form the hyperplane given as the kernel of \mathbf{v} and $(\mathbf{v}^T \beta) \mathbf{a}_0 = \beta_E \mathbf{a}_0$ is the intersection of $-\text{qdeg}_A(S_{A'}) + \beta$ with $\mathbb{C}\mathbf{a}_0$.

If $\mathbf{a}_0 \in \mathbb{R}_{\geq 0}A'$ then $\mathbb{N}\mathbf{a}_0$ meets $\mathbb{N}A'$ and so $\partial_0^k = \partial^{\mathbf{u}}$ with $\mathbf{u} = (0, u_1, \dots, u_n) \in \mathbb{N}A'$. In particular, $J_{A,0} = S_{A'}$ in this case. Moreover, $(x_0 \partial_0)(x_0 \partial_0 - 1) \cdots (x_0 \partial_0 - k + 1) = x_0^k \partial_0^k = x_0^k (\partial_0^k - \partial^{\mathbf{u}}) + x_0^k \partial^{\mathbf{u}} \in H_A(\beta) + V^1(D_A)$ shows the claim made in this case.

Now suppose that A and A' have equal rank but $\mathbf{a}_0 \notin \mathbb{R}_{\geq 0}A'$. In that case, $\bar{J}_{A,0}$ is a non-trivial ideal of $S_{A'}$. We shall use a toric filtration

$$(N) \quad : \quad 0 = N_0 \subsetneq N_1 \subsetneq \dots \subsetneq N_t = S_{A'}/\bar{J}_{A,0}$$

and let $J_\alpha \supseteq J_{A,0}$ be the $R_{A'}$ -ideal such that $N_\alpha = J_\alpha/J_{A,0}$. We will view J_α as subset of $D_{A'}$ or even D_A . In analogy to the previous case, for any $\partial^{\mathbf{u}}$ in $J_{A,0;k}$ the b -function along x_0 of the coset of $\partial^{\mathbf{u}}$ in $M_A(\beta)$ divides $s(s-1) \cdots (s-k+1)$. Indeed, $\partial^{\mathbf{u}} \in J_{A,0;k}$ implies that $\partial_0^k \partial^{\mathbf{u}} - \partial^{\mathbf{v}} \in I_A$ for some \mathbf{v} with $v_0 = 0$, and so $x_0^k \partial_0^k \partial^{\mathbf{u}} \in H_A(\beta) + V^1(D_A)$. In particular, the root set of the b -function of the coset of $\partial^{\mathbf{u}}$ in $M_{A'}(\beta)$ is inside the set of integers described in the statement of the theorem.

For each composition factor $\bar{N}_\alpha = N_\alpha/N_{\alpha-1}$ choose now a facet τ_α of A' and an element $\partial^{\mathbf{u}_\alpha}$ of $S_{A'}$ $\mathbf{u}_\alpha \in \{0\} \times \mathbb{N}^n$ such that N_α is a quotient of $S_{A'} \cdot \partial^{\mathbf{u}_\alpha}$ and such that the annihilator of $\partial^{\mathbf{u}_\alpha}$ in \bar{N}_α contains the toric ideal $I_{A'}^{\tau_\alpha}$. Then $\text{qdeg}_{A'}(\bar{N}_\alpha)$ is contained in $A' \cdot \mathbf{u}_\alpha + \text{qdeg}_{A'}(S_{\tau_\alpha})$.

Since \mathbf{a}_0 is not in $\mathbb{R}_{\geq 0}A'$, Lemma 4.3 shows that the facet τ_α can be chosen such that $\mathbf{a}_0 \notin \mathbb{Q} \cdot \tau_\alpha$. Indeed, if an entire face of $\mathbb{R}_{\geq 0}A'$ is visible from \mathbf{a}_0 then it sits in at least one facet whose span does not contain \mathbf{a}_0 . By Lemma 2.3 there is an element E_α of the Euler space of A that does not involve any element of τ_α , but which has coefficient 1 for θ_0 . Notation 2.2 then associates a degree function $\text{deg}_{E_\alpha}(-)$ to α .

As $\partial_j \cdot \partial^{\mathbf{u}_\alpha} \in N_{\alpha-1}$ for $j \notin \tau_\alpha$ it follows that the difference of $(E_\alpha - \beta_\alpha) \cdot \partial^{\mathbf{u}_\alpha}$ and $(\theta_0 - \beta_\alpha) \cdot \partial^{\mathbf{u}_\alpha}$ is inside $(V^0 D_A)N_{\alpha-1}$. Since $E_\alpha - \beta_\alpha$ is in $H_A(\beta)$, so is $\partial^{\mathbf{u}_\alpha}(E_\alpha - \beta_\alpha) = (E_\alpha - \beta_\alpha + \text{deg}_{E_\alpha}(\partial^{\mathbf{u}_\alpha}))\partial^{\mathbf{u}_\alpha}$. Therefore, $(\theta_0 - \beta_\alpha + \text{deg}_{E_\alpha}(\partial^{\mathbf{u}_\alpha}))\partial^{\mathbf{u}_\alpha}$ is in $H_A(\beta) + (V^0 D_A)N_{\alpha-1}$. Then, in parallel to how Lemma 3.4 was used in the proof of Theorem 3.5, the product

$$\prod_{\alpha} (\theta_0 - \beta_\alpha + \text{deg}_{E_\alpha}(\partial^{\mathbf{u}_\alpha}))$$

multiplies $1 \in D_A$ into $H_A(\beta) + (V^0 D_A)J_{A,0} + (V^1 D_A)$. Multiplying by $x_0^k \partial_0^k$ for suitable k one obtains the desired bound for the b -function as in the second paragraph of the proof.

It follows as in Theorem 3.5 (with the modification that we have here θ_0 rather than $\mathcal{F}^{-1}(\theta_0)$, which affects signs) that the intersection of the roots of all such bounds is the intersection of $(-q\deg_{A'}(S_{A'}/\bar{J}_{A,0}) + \beta)$ with the line $\mathbb{C} \cdot \mathbf{a}_0$. ■

Example 4.5. With $A = (\mathbf{a}_0, \mathbf{a}_1, \mathbf{a}_2) = \begin{pmatrix} -1 & 0 & 3 \\ 1 & 1 & 1 \end{pmatrix}$, consider the b -function along x_1 of the A -hypergeometric system. The ideal $J_{A,1}$ is generated by $1 \in S_{A'} = \mathbb{C}[\mathbb{N}(\mathbf{a}_0, \mathbf{a}_2)]$ since ∂_1^4 is in $S_{A'}$. The set of necessary integer roots is then $\{0, 1, 2, 3\}$. No other roots are needed since $S_A/J_{A,1}$ is zero, irrespective of β . Figure 3 shows the situation in this case.

Restriction to $(x_2 = 0)$ behaves differently. As $S_{A'} = \mathbb{C}[\mathbb{N}(\mathbf{a}_0, \mathbf{a}_1)]$ now, $J_{A,2} = J_{A,2;1}$ is generated by ∂_0^3 , and the quasi-degrees of $S_{A'}/J_{A,2}$ are the lines $\mathbb{C} \cdot (0, 1) + (i, 0)$ with $i = 0, -1, -2$. The intersection of the negative of these three lines, shifted by β , with the line $\mathbb{C} \cdot \mathbf{a}_2$ is $\mathbf{a}_2 \cdot \{(i + \beta_1)/3\}_{i=0,1,2}$. So the b -function has (at worst) roots $\{0, \beta_1, \beta_1 + 1, \beta_1 + 2\}/3$. Figure 4 shows the quasi-degrees of $S_A/J_{A,2}$. □




Fig. 3. The elements of $S_A \setminus S_{A'}$ (black) and $S_{A'}$ (shaded) for restriction to x_1 .




Fig. 4. The quasi-degrees of $S_A/J_{A,2}$ form three parallel lines.

Remark 4.6. We believe that both bounds in Theorems 3.5 (as is) and 4.4 (up to integers) are sharp. \square

4.2 Restriction to a generic point

We suppose here that A is homogeneous; in other words, the Euler space contains a homothety. Let $p = (p_0, \dots, p_n)$ be a point of \mathbb{C}^{n+1} . We wish to estimate here the b -function for restriction of $M_A(\beta)$ to the point $-p$ if p is generic. As a holonomic module is a connection near any generic point, this restriction yields a vector space isomorphic to the space of solutions to $H_A(\beta)$ near $-p$, see [12, Section 5.2].

Definition 4.7. Let $\theta_p = (x_0 + p_0)\partial_0 + \dots + (x_n + p_n)\partial_n$ and write θ for θ_p if $p = 0$. The b -function for restriction of a principal D -module $M = D/I$ to the point $x + p = 0$ is the minimal polynomial $b_p(s)$ such that $b_p(\theta_p) \in I + (V_p^1 D)$ where $V_p^k D$ is the Kashiwara–Malgrange V -filtration along $\text{Var}(x + p)$:

$$V_p^k D = \mathbb{C} \cdot \{(x + p)^{\mathbf{u}} \partial^{\mathbf{v}} \mid |\mathbf{u}| - |\mathbf{v}| \geq k\}. \quad \square$$

Remark 4.8.

1. For any pair of manifolds $Y \subseteq X$ and given a D -module M on X one can define a b -function of restriction for the section $m \in M$ along Y by a formula generalizing both Definitions 1.1 and 4.7. Kashiwara proved their existence for holonomic M .
2. The roots of this b -function here relate to restriction of solution sheaves as follows. Near a generic point $x + p = 0$, a D -module M is a connection whose solution space has a basis consisting of a certain number of holomorphic functions. The germs of these functions form a vector space that can be identified with the dual of the zeroth homology group of $(D/(x + p)D) \otimes_D^L M$. Filtering this complex by $V_p^\bullet D$, $b_p(k)$ annihilates the k -th graded part of its homology, compare [7, 8, 16]. In particular, $b_p(s)$ carries information on the starting terms of the solution sheaf of M near $x + p = 0$. \square

The purpose of this section is to bound $b_p(s)$ for $I = H_A(\beta)$ and generic p with the following strategy. We first show that a polynomial $b(s)$ is a multiple of $b_p(s)$ if $b(\theta)$

is in $D_A(I_A, A \cdot \mathcal{E} \cdot \partial)$ where

$$\mathcal{E} = \begin{pmatrix} p_0 & 0 & \cdots & 0 \\ 0 & p_1 & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \cdots & 0 & p_n \end{pmatrix},$$

provided that p is component-wise non-zero. The generators of $D_A(I_A, A \cdot \mathcal{E} \cdot \partial)$ are independent of x and we next observe that the radical of $R_A(I_A, A \cdot \mathcal{E} \cdot \partial)$ is $R_A \cdot \partial$, provided that p is generic. Thus, $b_p(s)$ will be a factor of any polynomial that annihilates the finite length module $R_A/(I_A, A \cdot \mathcal{E} \cdot \partial)$ as long as p is generic. We exhibit a particular such polynomial with all roots integral. In the case of a normal semigroup ring, we show that the (necessarily integral) roots of $b_p(s)$ are in the interval $[0, d - 1]$.

We begin with pointing out that $b(\theta_p) \in I + (V_p^1 D)$ is equivalent to $b(\theta) \in I_p + (V_0^1 D)$, where I_p is the image of I under the morphism induced by $x \mapsto x - p$, $\partial \mapsto \partial$ and $(V_0^k D)$ is the Kashiwara–Malgrange filtration along the origin. Among the generators of $I = H_A(\beta)$, only the Euler operators depend on x while $(I_A)_p = I_A$ for any p ; one has $(E_i - \beta_i)_p = \sum a_{i,j}(x_j - p_j)\partial_j - \beta_i = E_i - \beta_i - \sum a_{i,j}p_j\partial_j$. We hence seek a relation $b(\theta) \in D_A \cdot (I_A, E - \beta - A \cdot \mathcal{E} \cdot \partial) + (V_0^1 D_A)$ with \mathcal{E} as in the display above.

Generally, a statement $b(\theta) \in I + (V_0^1 D_A)$ is equivalent to $b(\theta)$ being in the degree zero part $\text{gr}_{V_0}^0(I)$ of the associated graded object. Note that $\text{gr}_{V_0}(D_A)$ is a Weyl algebra again (although of course the symbol map $D_A \rightarrow \text{gr}_{V_0}(D_A)$ is not an isomorphism). Abusing notation, we denote x and ∂ also the symbols in $\text{gr}_{V_0}(D_A)$ of the respective elements of D_A . By the previous paragraph then, the graded ideal $\text{gr}_{V_0}(H_A(\beta)_p)$ contains the elements that generate I_A (since I_A is homogeneous!), as well as the elements $A \cdot \mathcal{E} \cdot \partial$ which arise as the V_0 -symbols of $E_p - \beta$.

We need the following folklore result) for which we know no explicit reference.

Claim. The R_A -ideal generated by I_A and $A \cdot \mathcal{E} \cdot \partial$ has, for generic \mathcal{E} , radical $R_A \cdot \partial$. \square

A sequence of d generic linear forms is of course a system of parameters on S_A ; the issue is to show that linear forms of the type $A \cdot \mathcal{E} \cdot \partial$ are sufficiently generic.

Proof. As I_A and $A \cdot \mathcal{E} \cdot \partial$ are standard graded, $\text{Var}(I_A, A \cdot \mathcal{E} \cdot \partial)$ is a conical variety. It thus suffices to show that the ideal $\text{Var}(I_A, A \cdot \mathcal{E} \cdot \partial)$ is of height $n + 1$.

The ideal $R_A[x](I_A, A \cdot \theta)$ in the polynomial ring $R_A[x]$ defines in the cotangent bundle $\text{Spec}(R_A[x])$ of \mathbb{C}^{n+1} the union of the conormals to each torus orbit since the

Euler fields are tangent to the torus and span a space of the correct dimension in each orbit point. Suppose the claim is false, so that there is a non-zero point $y \in \text{Var}(I_A)$ such that (the generically chosen vector) p is a conormal vector to the orbit of y . If y is in a torus orbit O_τ associated to a proper face τ of A then its coordinates corresponding to $A \setminus \tau$ are zero and we can reduce the question to the case where $A = \tau$. It is hence enough to show that there is $p \in \mathbb{C}^{n+1}$ such that p is not a conormal vector to any smooth point of $\text{Var}(I_A)$.

Let $X \subseteq \mathbb{C}^{n+1}$ be any reduced affine variety and denote X_0 its smooth locus. We define a set $C(X)$ inside \mathbb{C}^{n+1} by setting

$$[\eta \in C(X)] \iff [\exists y \in X_0, \quad \eta \in (T_{X_0}^*(\mathbb{C}^{n+1}))_y]$$

where $(T_{X_0}^*(\mathbb{C}^{n+1}))_y$ is the fiber of the conormal bundle at y of the pair $X_0 \subseteq \mathbb{C}^{n+1}$. This is a constructible, analytically parameterized union of a $\dim(X)$ -dimensional family of vector spaces of dimension $n + 1 - \dim(X)$, which hence might fill \mathbb{C}^{n+1} .

Now suppose that X is a conical variety; then the conormals of y and λy agree for all $\lambda \in \mathbb{C}^*$. In particular,

$$C(X) = \bigcup_{\bar{y} \in \text{Proj}(X)} (T_{X_0}^*(\mathbb{C}^{n+1}))_{\bar{y}},$$

where $\text{Proj}(X)$ is the associated projective variety. But this is now an analytically parameterized union of a $(\dim(X) - 1)$ -dimensional family of vector spaces of dimension $n + 1 - \dim(X)$. It follows that most elements of \mathbb{C}^{n+1} are outside $C(X)$ in this case, and the claim follows. \blacksquare

It follows from the Claim that $\text{gr}_{V_0}(H_A(\beta)_p)$ contains all monomials in ∂ of a certain degree k that depends on A . Let $E = \theta_0 + \dots + \theta_n$; by hypothesis $E - \beta_E \in H_A(\beta)$.

Lemma 4.9. Denote ∂_A^k the set of all monomials of degree k in $\partial_0, \dots, \partial_n$, and $D_A \cdot \partial_A^k$ the left D_A -ideal generated by ∂_A^k . Then in $D_A/D_A \cdot \partial_A^k$, the identity $E(E - 1) \dots (E - k + 1) \cong 0$ holds. \square

Proof. This is clear if $k = 1$. In general, by induction,

$$E(E - 1) \dots (E - k + 1) \in D_A \cdot \partial_A^{k-1} \cdot (E - k + 1) = D_A \cdot E \cdot \partial_A^{k-1} \subseteq D_A \cdot \partial_A^k. \quad \blacksquare$$

Remark 4.10. The homogeneity of X is necessary in the Claim, since otherwise $C(X)$ does not need to be contained in a hypersurface. Consider, for example, $A = (2, 1)$ in

which case the union of all tangent lines (nearly) fills the plane, and where the zero locus of I_A and $A \cdot \mathcal{E} \cdot \partial$ contains always at least two points. \square

The lemma implies that $\text{gr}_{V_0}^0(H_A(\beta)_p)$ contains $E(E-1) \dots (E-k+1)$ if p is generic. In other words, the b -function for restriction of $M_A(\beta)$ to a generic point divides $s(s-1) \dots (s-k+1)$.

In some cases, one can be more explicit about $k-1$, the top degree in which $R_A/R_A(I_A, A \cdot \mathcal{E} \cdot \partial)$ is non-zero. Suppose S_A is a Cohen–Macaulay ring, then systems of parameters are regular sequences. In particular, the Hilbert series of $Q_A := R_A/R_A(I_A, A \cdot \mathcal{E} \cdot \partial)$ is that of S_A multiplied by $(1-t)^d$. Suppose in addition, that S_A is normal. Since, we already assume that S_A is standard graded, let P be the polytope that forms the convex hull of the columns of A . The Hilbert series of S_A is then of the form $\sum_{m=0}^{\infty} p_m \cdot t^m$, where p_m is the number of lattice points in the dilated polytope $m \cdot P$. This number of lattice points is counted by the Ehrhart polynomial $E_P(m)$ of P , a polynomial of degree $d-1 = \dim(P)$. If one writes the Hilbert series of S_A in standard form $Q(t)/(1-t)^d$ then the Hilbert series of Q_A is just the polynomial $Q(t)$. In particular, the highest degree of a non-vanishing element of Q_A is the degree of $Q(t)$.

In order to determine $\deg(Q(t))$ let $E_P(m) = e_{d-1}m^{d-1} + \dots + e_0$. Now in

$$\sum_{m=0}^{\infty} E_P(m)t^m = \sum_{i=0}^{d-1} \left(e_i \cdot \sum_{m=0}^{\infty} m^i \cdot t^m \right),$$

each term $\sum_{m=0}^{\infty} m^i \cdot t^m$, for $m > 0$, is a polylogarithm $\text{Li}_{-i}(t)$ given by $(t \frac{d}{dt})^i (\frac{t}{1-t})$. A simple calculation shows that $\text{Li}_{-i}(t)$ is the quotient of a polynomial of degree $i-1$ by $(1-t)^i$. Hence the sum in the display is the quotient of a polynomial of degree at most $d-1$ by $(1-t)^d$. The degree is truly $d-1$ as one can check from the differential expression for $\text{Li}_{-i}(t)$.

Therefore, the Hilbert series $Q(t)$ of Q_A is a polynomial of degree $d-1$. We have proved

Theorem 4.11. Let S_A be standard graded. The b -function for restriction of $M_A(\beta)$ to a generic point $x+p=0$ divides $s(s-1) \dots (s-k+1)$ where k denotes the highest degree in which the quotient $S_A/S_A \cdot (A \cdot \mathcal{E} \cdot \partial)$ is non-zero. If, in addition, S_A is normal then one may take $k=d$.

Funding

This work was supported by a DFG Emmy-Noether-Fellowship [RE 3567/1-1 to T.R.]; partially supported by the ANR/DFG joint program SISYPH [ANR-13-IS01-0001-01 and SE 1115/5-1 to T.R., C.S.]; and National Science Foundation [1401392-DMS to U.W.].

Acknowledgments

All three authors would like to thank the Forschungsinstitut Oberwolfach for hosting them in April of 2015. We are greatly indebted to an unknown referee for very careful reading, pointing out a number of misprints. □

References

- [1] Fernández-Fernández, M.-C. and U. Walther “Restriction of hypergeometric \mathcal{D} -modules with respect to coordinate subspaces.” *Proceedings of the American Mathematical Society* 139, no. 9 (2011): 3175–80.
- [2] Gel’fand, I. M. “General theory of hypergeometric functions.” *Doklady Akademii Nauk SSSR* 288, no. 1 (1986): 14–18.
- [3] Hotta, R. Equivariant D -modules. *arXiv:math/9805021*, pages 1–30, 1998.
- [4] Kashiwara, M. “ B -functions and holonomic systems. Rationality of roots of B -functions.” *Inventiones mathematicae* 38, no. 1 (1976/77): 33–53.
- [5] Maisonobe, P. and Z. Mebkhout. “Le théorème de comparaison pour les cycles évanescents.” In *Éléments De La Théorie Des Systèmes Différentiels Géométriques*, volume 8 of Sémin. Congr., 311–89. France: Soc. Math., 2004.
- [6] Matusевич, L. F., E. Miller, and U. Walther. “Homological methods for hypergeometric families.” *Journal of the American Mathematical Society* 18, no. 4 (2005): 919–41 (electronic).
- [7] Oaku, T. “Algorithms for b -functions, restrictions, and algebraic local cohomology groups of D -modules.” *Advances in Applied Mathematics* 19, no. 1 (1997): 61–105.
- [8] Oaku, T. and N. Takayama. “Algorithms for D -modules—restriction, tensor product, localization, and local cohomology groups.” *Journal of Pure and Applied Algebra* 156, no. (2–3) (2001): 267–308.
- [9] Reichelt, T. “Laurent polynomials, GKZ-hypergeometric systems and mixed Hodge modules.” *Compositio Mathematica* 150, no. 6 (2014): 911–41.
- [10] Reichelt, T. and C. Sevenheck. “Hypergeometric Hodge modules.” *arXiv:1503.01004*, pages 1–48, 2015.
- [11] Saito, M. “Modules de Hodge polarisables.” *Publications of the Research Institute for Mathematical Sciences* 24, no. 6 (1989): 849–995, 1988.
- [12] Saito, M., B. Sturmfels, and N. Takayama. *Gröbner Deformations of Hypergeometric Differential Equations*, volume 6 of Algorithms and Computation in Mathematics. Berlin: Springer, 2000.

- [13] Schulze, M. and U. Walther. "Irregularity of hypergeometric systems via slopes along coordinate subspaces." *Duke Mathematical Journal* 142, no. 3 (2008): 465–509.
- [14] Schulze, M. and U. Walther. "Cohen-Macaulayness and computation of Newton graded toric rings." *Journal of Pure and Applied Algebra* 213, no. 8 (2009): 1522–35.
- [15] Schulze, M. and U. Walther. "Hypergeometric D-modules and twisted Gauß-Manin systems." *Journal of Algebra* 322, no. 9 (2009): 3392–9.
- [16] Walther, U. "Algorithmic computation of de Rham cohomology of complements of complex affine varieties." *Journal of Symbolic Computation* 29, no. (4–5) (2000): 795–839. Symbolic computation in algebra, analysis, and geometry (Berkeley, CA, 1998).